

Supplementary Information for

Bioinformatics-Guided Discovery of miRNAs Involved in Apoptosis Modulated by Parthenolide Combined with Vincristine in The NALM6 Cell Line

Atefeh Bahmei, M.Sc.¹, Sepideh Namdari, M.Sc.², Mohammad Yaghoubzad-Maleki, M.Sc.³, Ali Emami, M.Sc.⁴,
Reza Ranjbaran, Ph.D.², Gholamhossein Tamaddon, Ph.D.^{1, 2*} 

1. Division of Hematology and Blood bank, Department of Medical Laboratory Sciences, School of Paramedical Sciences, Shiraz university of medical Sciences, Shiraz, Iran
2. Diagnostic Laboratory Sciences and Technology Research Center, School of Paramedical Sciences, Shiraz University of Medical Sciences, Shiraz, Iran
3. Division of Biochemistry, Department of Animal Biology, Faculty of Natural Sciences, University of Tabriz, Iran
4. Department of Biochemistry and Molecular Medicine, University of Montreal, Montreal, Canada




Fig.S1: The flow cytometry analysis following different treatments in NALM6 cells after 48 hours. **A.** Untreated cells, **B.** Cells treated with 0.8 nM vincristine, **C.** 1.2 nM vincristine, **D.** 1.925 μ M parthenolide, **E.** 2.8 μ M parthenolide, and **F.** 1.925 μ M parthenolide combined with 1.2 nM vincristine.

Received: 16/October/2023, Revised: 10/January/2024, Accepted: 21/January/2024

*Corresponding Address: P.O.Box: 7134814336, Division of Hematology and Blood bank, Department of Medical Laboratory Sciences, School of Paramedical Sciences, Shiraz university of medical Sciences, Shiraz, Iran
Email: tamaddon@sums.ac.ir