

Supplementary Information for

Betaine Attenuates The Expression of Vasoactive Mediators and Histological Alterations Associated with Ovarian Hyperstimulation Syndrome in Rats

Mohammad Reza Abbasi, D.V.M.¹, Javad Jamshidian, Ph.D.², Zahra Basir, Ph.D.^{3*} D, Mohammad Reza Tabandeh, Ph.D.4,5

- 1. Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran

- 2. Department of Pharmacology, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran
 3. Department of Basic Sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran
 4. Department of Biochemistry and Molecular Biology, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran
 5. Stem Cells and Transgenic Technology Research Center, Shahid Chamran University of Ahvaz, Ahvaz, Iran

Table S1: Comparison of the weight changes in the study groups

Parameters	Control	OHSS	OHSS+BET	OHSS+CAB	BET	CAB
Weight delta (g)	83.2 ± 2.5	$132.1 \pm 10.8^*$	$104.4 \pm 4.2^{*\#}$	$88.3 \pm 2.8^{*\#}$	$85.9 \pm 2.8^{\#}$	84.8 ± 2.8#

Bilateral ANOVA analysis and Tukey's post hoc test were used for multiple comparisons. Results are written as mean ± SEM (n=5 per group). *; Significant change with respect to the control group (P<0.05), #; Significant change with respect to the OHSS group (P<0.05), OHSS; Ovarian hyperstimulation syndrome, BET; Betaine, and CAB; Cabergoline.

Received: 12/December/2022, Revised: 27/November/2023, Accepted: 23/ December/2023

*Corresponding Address: P.O.Box: 6135783151, Department of Basic Sciences, Faculty of Veterinary Medicine, Shahid Chamran University of

Ahvaz, Ahvaz, Iran Email: z.basir@scu.ac.ir

