

# Repression of TGF- $\beta$ Signaling in Breast Cancer Cells by miR-302/367 Cluster

Mona Ahmadelizadeh Khanehsar, M.Sc.<sup>1,2</sup>, Moslem Hoseinbeyki, M.Sc.<sup>1</sup>, Masoumeh Fakhri Taha, Ph.D.<sup>1</sup>,

Arash Javeri, M.D., Ph.D.<sup>1\*</sup>

1. Department of Stem Cells and Regenerative Medicine, Institute for Medical Biotechnology, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran
2. Department of Biology, Damghan Branch, Islamic Azad University, Damghan, Iran

\*Corresponding Address: P.O. Box 14965-161, Department of Stem Cells and Regenerative Medicine, Institute for Medical Biotechnology, National Institute of Genetic Engineering and Biotechnology (NIGEB), Pajooheh Blvd., Tehran, Iran  
Email: arashj@nigeb.ac.ir

Received: 7/July/2018, Accepted: 21/November/2018

## Abstract

**Objective:** Epigenetic alterations of the malignantly transformed cells have increasingly been regarded as an important event in the carcinogenic development. Induction of some miRNAs such as miR-302/367 cluster has been shown to induce reprogramming of breast cancer cells and exert a tumor suppressive role by induction of mesenchymal to epithelial transition, apoptosis and a lower proliferation rate. Here, we aimed to investigate the impact of miR-302/367 overexpression on transforming growth factor- $\beta$  (TGF- $\beta$ ) signaling and how this may contribute to tumor suppressive effects of miR-302/367 cluster.

**Materials and Methods:** In this experimental study, MDA-MB-231 and SK-BR-3 breast cancer cells were cultured and transfected with miR-302/367 expressing lentivector. The impact of miR-302/367 overexpression on several mediators of TGF- $\beta$  signaling and cell cycle was assessed by quantitative real-time polymerase chain reaction (qPCR) and flow cytometry.

**Results:** Ectopic expression of miR-302/367 cluster downregulated expression of some downstream elements of TGF- $\beta$  pathway in MDA-MB-231 and SK-BR-3 breast cancer cell lines. Overexpression of miR-302/367 cluster inhibited proliferation of the breast cancer cells by suppressing the S-phase of cell cycle which was in accordance with inhibition of TGF- $\beta$  pathway.

**Conclusion:** TGF- $\beta$  signaling is one of the key pathways in tumor progression and a general suppression of TGF- $\beta$  mediators by the pleiotropically acting miR-302/367 cluster may be one of the important reasons for its anti-tumor effects in breast cancer cells.

**Keywords:** Breast Cancer, miR-302/367, Reprogramming, Transforming Growth Factor-Beta

Cell Journal (Yakhteh), Vol 21, No 4, January-March (Winter) 2020, Pages: 444-450

**Citation:** Ahmadelizadeh Khanehsar M, Hoseinbeyki M, Fakhri Taha M, Javeri A. Repression of TGF- $\beta$  Signaling in breast cancer cells by miR-302/367 cluster. Cell J. 2020; 21(4): 444-450. doi: 10.22074/cellj.2020.6193.

## Introduction

Despite recent advancements in the treatment of breast cancer, it still remains one of the leading causes of cancer deaths among women (1). Therefore, development of new therapeutic approaches for breast cancer is of the utmost importance. Reprogramming of somatic cells and generation of induced pluripotent stem (iPS) cells by some transcription factors including OCT3/4, SOX2, NANOG, KLF4, LIN28 and MYC (2, 3) demonstrated that the cell fate can be manipulated *in vitro*. Reprogramming is a process accompanied by distinct alterations in chromatin and transcriptional programs.

MiRNAs constitute a class of 17-24 bp small non-coding RNAs, involved in regulation of different biological processes and cancer-related cellular activities such as apoptosis, proliferation and invasion (4, 5). MiR-302/367 cluster possesses a coding sequence located in intron 8 of the *LARP7* gene and codes for 5 miRNAs including miR302a, miR302b, miR302c, miR302d, and miR367 which are highly expressed in embryonic stem cells (6-8), but their expression decline rapidly after differentiation (9). It was shown that miR-302/367 cluster can effectively

reprogram human and mouse somatic cells to iPS cells (10, 11). miR-302 is also able to reprogram human cancer cells to a human embryonic stem cell-like state with a slow cell cycle rate and dormant cell-like morphology (12, 13). Reprogramming by miR-302/367 cluster has shown tumor suppressive effects on different cancer cells, such as melanoma and colon cancer cells (14), cervical carcinoma cells (15) glioblastoma cells (16), prostate cancer cells (13), endometrial cancer cells (17) and breast cancer (18). The miR-302/367 cluster has been shown to induce reprogramming of somatic cells through multiple pathways, including MECP1/2 and AOF1/2 silencing, repression of suppressor *NR2F2* gene expression, and silencing RHOC and TGFBR2 (19).

Transforming growth factor- $\beta$  (TGF- $\beta$ ) signaling pathway is one of the major players in malignant progression through multiple mechanisms which enhance tumor cell invasion, dissemination, and immune evasion (20, 21). In this study we aimed to investigate how overexpression of miR-302/367 cluster in breast cancer cells affects some of the main TGF- $\beta$  signaling pathway mediators.

## Materials and Methods

### Cell lines and culture conditions

In this experimental study, human MDA-MB-231 and SK-BR-3 breast cancer cell lines were respectively purchased from Pasteur Institute and Iranian Biological Resource Center (IRBC), Iran. Both cell lines were cultured in Dulbecco's Modified Eagle's medium (DMEM) with 10% fetal bovine serum (FBS), 1% L-glutamine and 1% penicillin-streptomycin (all from Gibco™, Thermo Fisher Scientific, USA) at 5% CO<sub>2</sub> and 37°C. The culture medium was renewed every other day.

### Transfection with miR-302/367 expressing vector

Transfection of MDA-MB-231 and SK-BR-3 were performed using either a TDH101PA-GP miR-302abcd/367 expressing Lentivector (System Biosciences, SBI, USA) or the same vector without the miR-302/367 cluster as the mock control type, using Lipofectamine® 2000 transfection reagent (Invitrogen, Thermo Fisher Scientific, USA) according to the manufacturer's protocol. 48 hours after transfection, transfected cells were selected by adding 1 mg/ml puromycin dihydrochloride (Bio Basic Inc., Canada) to the culture medium every other day up to the elimination of untransfected cells. Transfected cells were kept in culture condition for a two-week period.

### Analysis of miRNA and gene expression by quantitative real time polymerase chain reaction

For analysis of miRNA expression, total RNA including

small RNA, was extracted from the cultured cells using RNX-Plus solution (Sinaclon, Iran) according to the manufacturer's protocol. Equal amounts of RNA were reverse transcribed into cDNA using BON-miR miRNA 1st-Strand cDNA Synthesis Kit (Stem Cell Technology Co., Iran).

For quantification of mRNAs, total RNA was extracted using the High Pure RNA Isolation Kit (Roche, Germany) according to the manufacturer's protocol. RNA quality and quantity were assessed using a NanoDrop™ 2000/2000c Spectrophotometer (Thermo Fisher Scientific, USA). Equal amount of total RNA from each group was reverse transcribed into cDNA using oligo-dT primers and RevertAid H Minus Reverse Transcriptase (Thermo Fisher Scientific). Assessment of miRNA and mRNA expression was performed, using FastStart SYBR Green Master (Roche, Germany) and specific primers for *miR-302a*, *miR-302b*, *miR-302c*, *miR-302d*, *miR-367* and other genes as mentioned in Table 1, on a Rotor-Gene 6000 (Corbett Research, Australia) real-time PCR instrument. *SNORD47* was selected as the internal reference gene for quantification of miRNAs. *GAPDH* and *B2M* were used as the internal reference genes for quantification of the mRNAs. Comparative analysis of gene expression between different groups was performed using REST 2009 software (Relative Expression Software Tool, Qiagen) based on a Pair Wise Reallocation Randomization Test (22). Four replicates of each group were included in the qPCR reactions.

**Table 1:** Primers used for quantitative real-time polymerase chain reaction

Target	Primer sequence (5'-3')	Size (bp)	Accession no.
<i>TGFBR2</i>	F: CCCATCCACTGAGACATATTAAT R: CATTCTTTCTCCATACAGCCAC	198	NM_001024847.2
<i>BUB1</i>	F: GAGTCAAATATGGAACGAAGAG R: GTCTTCATTTACCCATTGCTCA	207	NM_004336.4
<i>RHOC</i>	F: CCTGACAGCCTGGAAAACAT R: AACGGGCTCCTGCTTCATCT	153	NM_175744.4
<i>AKT1</i>	F: ACAAACGAGGGGAGTACATCAA R: TCTTCATCAGCTGGCACTGC	156	NM_005163.2
<i>MAPK1</i>	F: ATTCCAAGGGCTACACCAAGT R: GGATCCAAGAATACCCAAAATGT	136	NM_002745.4
<i>MAPK14</i>	F: TGGCTGTCGACTTGCTGGA R: CATAGGTCAGGCTTTTCCACT	189	NM_001315.2
<i>SMAD3</i>	F: CATAATAACTTGGACCTGCAGC R: ACGCCTCTTCCGATGTGTCT	236	NM_005902.3
<i>B2M</i>	F: TCCAGCGTACTCCAAAGATTCA R: GTCAACTTCAATGTCGGATGGAT	113	NM_004048.2
<i>GAPDH</i>	F: TCACCATCTTCCAGGAGCGA R: CAAATGAGCCCCAGCCTTCT	116	NM_002046.5

### Analysis of cell cycle by flow cytometry

At the end of transfection and cell culture period, the cells were harvested and fixed in 70% cold ethanol and DNA content was stained with propidium iodide (PI) solution. Four replicates of each group were used in this study. Cell cycle analysis was carried out using a FACSCalibur™ flow cytometer (BD Biosciences, USA). FlowJo vX.0.7 software (Tree Star Inc., USA) was used for analysis of the results. Comparison of the cell cycle G1, S and G2/M proportions was performed between the mock and miR-302/367 transfected group of each cell line, using unpaired t test.

## Results

### Overexpression of the miR-302/367 members in transfected cells

Antibiotic-based selection of the miR-302/367 transfected breast cancer cells caused producing a highly (>90%) GFP-expressing cells population (Fig.1A) which were used for the subsequent experiments. Quantification of the miR-302/367 expression in MDA-MB-231 cells showed upregulation of miR-302a, miR-302b, miR-302c, miR-302d and miR-367 by mean factors of 74, 946, 33, 145 and 25, respectively (Fig.1B). In SK-BR-3 cells, after miR-302/367 transfection, miR-302a, miR-302b, miR-


302c, miR-302d and miR-367 were upregulated by mean factors of 145, 1581, 20, 202 and 6, respectively (Fig.1B).

### Regulation of TGF-β and MAPK pathway genes by miR-302/367 cluster


Firstly, we checked how transfection of the breast cancer cells with miR-302/367 cluster affects the expression of some key mediators of TGF-β and mitogen-activated protein kinase (MAPK) pathways at gene level. Quantitative real-time PCR showed that in the both MDA-MB-231 and SK-BR-3 cells, overexpression of miR-302/367 cluster downregulated *TGFBR2*, *BUB1*, *RHOC*, *AKT1*, *MAPK1*, *MAPK14* and *SMAD3* expression compared to the mock transfected cells (Fig.2).

### Cell cycle arrest by overexpression of miR-302/367 cluster


At the end of culture period, transfected breast cancer cells with either miR-302/367 or mock vector were analyzed for the cell cycle phases, using PI staining and flow cytometry. In the miR-302/367 transfected MDA-MB-231 and SK-BR-3 cells, there was a marked decrease in the S-phase population, while the G2/M phase population was partially increased compared to the mock transfected group (Fig.3).



**Fig.1:** Ectopic expression of miR-302 cluster in the BC cells. **A.** Photomicrographs of the MDA-MB-231 and SK-BR-3 cells transfected with either miR-302/367 or mock vector. Transfected cells show GFP expression. Scale bar represents 50 μm and **B.** Assessment of miR-302/367 expression using quantitative real-time polymerase chain reaction (qPCR) in MDA-MB-231 cells (left) and SK-BR-3 cells (right) transfected with miR-302/367 vector. Fold changes are reflected on the vertical axis compared to the control group (transfected with mock vector) which has been normalized to 1. Analysis performed by REST 2009 software based on a Pair Wise Fixed Reallocation Randomisation Test© and significant P values (\*, P<0.05, \*\*\*, P<0.001) are indicated on the chart. BC; Breast cancer and GFP; Green fluorescent protein.




**Fig.2:** Expression analysis of some transforming growth factor-beta (TGF-β) mediators at mRNA level after transfection with miR-302/367 vector using quantitative real-time polymerase chain reaction (qPCR). Downregulation of TGF-β-related genes in **A.** MDA-MB-231 and **B.** SK-BR-3 cells. Red line represents expression level in the mock transfected group. P values were generated by REST 2009 software based on a Pair Wise Fixed Reallocation Randomisation Test©. Significant P values (\*; P<0.05, \*\*; P<0.01, \*\*\*; P<0.001) are reflected on the chart.



**Fig.3:** Flow cytometry analysis of cell cycle. There was a significant decrease in S-phase and a partial increase in G2/M-phase population of both MDA-MB-231 and SK-BR-3 cells, after overexpression of miR-302/367 cluster (unpaired t test, n=4, \*, P<0.05, \*\*, P<0.01 and \*\*\*, P<0.001).





**Fig.4:** Interaction between miR-302/367 cluster and several mediators of transforming growth factor-beta (TGF-β) signaling in both canonical and non-canonical pathways. The inhibitory effect of miR-302/367 cluster is shown by the red lines.

## Discussion

Genetic and epigenetic alterations contribute to cancer initiation and progression through affecting gene expression. While genetic mutations may lead to stable and irreversible alterations, transient and reversible changes are usually caused by epigenetic modifications (23). It has been shown that reprogramming of cancer cells by some pluripotency transcription factors or specific miRNAs, like miR-302/367 cluster, may lead to an embryonic stem cell-like state and less tumorigenicity (13, 24). We previously demonstrated upregulation of some pluripotency factors, including OCT4A, SOX2 and NANOG, by overexpression of miR-302/367 cluster in MDA-MB-231 and SK-BR-3 cells (18).

Accumulating evidence supports the function of miR-302 cluster as a tumor suppressor family which can alleviate tumorigenicity of cancer cells through reversal of epithelial to mesenchymal transition (EMT), induction of apoptosis and anti-proliferative effect (13-15). Previously, we demonstrated some anti-tumor effects of miR-302 cluster in melanoma, colon and breast cancer cells (14, 18). Among the pathways promoting EMT, several studies have reported that inhibition of TGF-β signaling induces somatic cell reprogramming (25, 26).

In this study, we investigated how overexpression of miR-302/367 cluster in MDA-MB-231 and SK-BR-3 breast cancer cells affects some mediators of TGF-β/MAPK/AKT signaling pathway at gene level. As shown, *TGFBR2* and *RHOC* are directly targeted by miR-302 cluster, subsequently facilitating human or mouse fibroblast reprogramming towards iPS cells (27,

28). In accordance with these studies, we found that overexpression of miR-302/367 in human breast cancer cells downregulates *TGFBR2* and *RHOC* expressions. TGF-β signaling has two canonical and non-canonical pathways (Fig.4). In the canonical or SMAD-dependent pathway, SMAD proteins play key regulatory roles among which SMAD2 and SMAD3 proteins are phosphorylated through activity of TGF-β and activin (29). While, in the non-canonical or SMAD-independent pathway, TGF-β activates phosphatidylinositol 3kinase (PI3K)/AKT and MAPK pathways (30). In the current study, expressions of *TGFBR2*, *BUB1*, *RHOC*, *AKT1*, *MAPK1*, *MAPK14* and *SMAD3* were significantly downregulated after overexpression of miR-302/367 cluster in the both cell lines. Previously, Cai et al. (15) reported that miR-302/367 directly targets *AKT1* and it suppresses proliferation of HeLa and SiHa cervical carcinoma cells. In the same study, AKT1 protein level was decreased after miR-302/367 transfection, but *AKT1* gene expression was not significantly changed. In another study, Li et al. (31) demonstrated that miR-302abcd cluster upregulated OCT4 expression by targeting *AKT1* gene at its 3'-UTR. The same report also showed downregulation of AKT1 at both gene and protein levels, following miR-302 transfection. Similarly, we showed that overexpression of miR-302/367 cluster in MDA-MB-231 and SK-BR-3 cells induces expression of *OCT4* gene (18) and downregulates expression of *AKT1*. Therefore, it seems that a mechanism, similar to that of previous reports, is applicable to breast cancer cells.

We also detected a significant downregulation of *SMAD3* expression following miR-302/367 transfection

of the breast cancer cells. Sustained activity of SMAD complexes in the nucleus is one of the key features of TGF- $\beta$  signaling in cancer cells. It was reported that an interaction between FOXM1 and SMAD3 is critical for TGF- $\beta$ -mediated gene expression. Thus, it promotes breast cancer cell invasion and metastasis (32).

*BUB1* was also downregulated in the breast cancer cells after ectopic expression of *miR-302/367* cluster. BUB1 is a serine/threonine kinase playing a significant role in cell cycle regulation, chromosome cohesion (33), and it is a key mediator of TGF- $\beta$  signaling. It has been shown that BUB1 promotes canonical and non-canonical TGF- $\beta$  signaling and mediates TGF- $\beta$ -dependent EMT, cell migration and invasion through interaction with both TGFBR1 and TGFBR2 (34). Here, for the first time, we are reporting downregulation of *BUB1* expression in breast cancer cells following overexpression of *miR-302/367* cluster. This provides further evidence regarding the significance of *miR-302/367* suppressive impact on TGF- $\beta$  signaling through inhibition of both canonical and non-canonical pathways.

MAPK pathway is part of the non-SMAD pathways, activated by the TGF- $\beta$  receptors (35). MAPK1, also known as ERK2, is encoded by *MAPK1* gene. The ERK1/2 pathway plays a pivotal role in regulation of cell proliferation, and it is known as a master regulator of G1 to S-phase progression (36, 37). Another player of the non-canonical TGF- $\beta$  pathway, MAPK14/p38 $\alpha$  is encoded by *MAPK14* gene. MAPK14/p38 $\alpha$  is 50% identical to ERK2 and generally expressed in cell lines and tissues (36). There has been controversy regarding the role of p38 MAPKs in regulating cell proliferation and survival (38). This primarily depends on the cell type determining whether p38 MAPK induces progression or inhibition of G1/S transition through differential regulation of cyclin A or D levels, phosphorylation of RB protein (39, 40), and phosphorylation of p53 (40). In our study, ectopic expression of *miR-302/367* cluster in the breast cancer cells downregulated expression of all of the investigated TGF- $\beta$  mediators, including *TGFBR2*, *BUB1*, *RHOC*, *AKT1*, *MAPK1*, *MAPK14* and *SMAD3*. These findings indicate a strong suppressive effect of *miR-302/367* cluster on the TGF- $\beta$  signaling (Fig.4). In this study we report a lower proliferation rate and S-phase suppression of the breast cancer cells by overexpression of *miR-302/367*, confirming our previous report (18). This can be explained by suppression of *MAPK1* and *MAPK14* to some extent. Therefore, suppression of TGF- $\beta$  mediators may provide a good reason behind the partial cell cycle arrest observed in the breast cancers, following overexpression of *miR-302/367* cluster.

## Conclusion

Overexpression of *miR-302/367* cluster in human breast cancer cells resulted in a general suppressive effect on multiple mediators of TGF- $\beta$  signaling and BUB1. This finding was accompanied by inhibition of cell proliferation. Previously, we reported anti-tumor

effects of either *miR-302bcad* or *miR-302bcad/367* clusters on melanoma, colon and breast cancer cells due to induction of apoptosis and suppression of proliferation and invasion. Current results are providing new evidence that suppression of TGF- $\beta$  signaling at gene level may be one of the important reasons for anti-tumor effects of *miR-302/367* cluster in breast cancer cells.

## Acknowledgements

There is no financial support and conflict of interest in this study.

## Authors' Contributions

M.A.K.; Did the gene expression analysis and writing the draft. M.H.; Performed cell culture and transfection of the cells. M.F.T.; Gave technical assistance on the reprogramming process and contributed to writing the draft. A.J.; Designed the study, supervised the project, performed the cell cycle analysis and edited the manuscript draft. All authors read and approved the final manuscript.

## References

- Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin*. 2015; 65(2): 87-108.
- Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*. 2006; 126(4): 663-676.
- Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, et al. Induced pluripotent stem cell lines derived from human somatic cells. *Science*. 2007; 318(5858): 1917-1920.
- Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*. 2004; 116(2): 281-297.
- Ma L, Weinberg RA. MicroRNAs in malignant progression. *Cell Cycle*. 2008; 7(5): 570-572.
- Pi J, Tao T, Zhuang T, Sun H, Chen X, Liu J, et al. A microRNA302-367-Erk1/2-Klf2-S1pr1 Pathway prevents tumor growth via restricting angiogenesis and improving vascular stability. *Circ Res*. 2017; 120(1): 85-98.
- Zhang Z, Hong Y, Xiang D, Zhu P, Wu E, Li W, et al. MicroRNA-302/367 cluster governs hESC self-renewal by dually regulating cell cycle and apoptosis pathways. *Stem Cell Reports*. 2015; 4(4): 645-657.
- Suh M-R, Lee Y, Kim JY, Kim S-K, Moon S-H, Lee JY, et al. Human embryonic stem cells express a unique set of microRNAs. *Dev Biol*. 2004; 270(2): 488-498.
- Kuo CH, Ying SY. Advances in microRNA-mediated reprogramming technology. *Stem Cells Int*. 2012; 2012: 823709.
- Anokye-Danso F, Trivedi CM, Juhr D, Gupta M, Cui Z, Tian Y, et al. Highly efficient miRNA-mediated reprogramming of mouse and human somatic cells to pluripotency. *Cell Stem Cell*. 2011; 8(4): 376-388.
- Galoian K, Qureshi A, D'Ippolito G, Schiller PC, Molinari M, Johnstone AL, et al. Epigenetic regulation of embryonic stem cell marker miR302C in human chondrosarcoma as determinant of anti-proliferative activity of proline-rich polypeptide 1. *Int J Oncol*. 2015; 47(2): 465-472.
- Lin SL, Chang DC, Ying SY, Leu D, Wu DT. MicroRNA miR-302 inhibits the tumorigenicity of human pluripotent stem cells by coordinate suppression of the CDK2 and CDK4/6 cell cycle pathways. *Cancer Res*. 2010; 70(22): 9473-9482.
- Lin SL, Chang DC, Chang-Lin S, Lin CH, Wu DT, Chen DT, et al. Mir-302 reprograms human skin cancer cells into a pluripotent ES-cell-like state. *RNA*. 2008; 14(10): 2115-2124.
- Maadi H, Moshtaghian A, Taha MF, Mowla SJ, Kazeroonian A, Haass NK, et al. Multimodal tumor suppression by miR-302 cluster in melanoma and colon cancer. *Int J Biochem Cell Biol*. 2016; 81(Pt A): 121-132.

15. Cai N, Wang YD, Zheng PS. The microRNA-302-367 cluster suppresses the proliferation of cervical carcinoma cells through the novel target AKT1. *RNA*. 2013; 19(1): 85-95.
16. Yang CM, Chiba T, Brill B, Delis N, von Manstein V, Vafaizadeh V, et al. Expression of the miR-302/367 cluster in glioblastoma cells suppresses tumorigenic gene expression patterns and abolishes transformation related phenotypes. *Int J Cancer*. 2015; 137(10): 2296-2309.
17. Yan GJ, Yu F, Wang B, Zhou HJ, Ge QY, Su J, et al. MicroRNA miR-302 inhibits the tumorigenicity of endometrial cancer cells by suppression of Cyclin D1 and CDK1. *Cancer Lett*. 2014; 345(1): 39-47.
18. Ramezankhani B, Taha MF, Javeri A. Vitamin C counteracts miR-302/367-induced reprogramming of human breast cancer cells and restores their invasive and proliferative capacity. *J Cell Physiol*. 2019; 234(3): 2672-2682.
19. Kuo CH, Deng JH, Deng Q, Ying SY. A novel role of miR-302/367 in reprogramming. *Biochem Biophys Res Commun*. 2012; 417(1): 11-16.
20. Miyazono K. Transforming growth factor-beta signaling in epithelial-mesenchymal transition and progression of cancer. *Proc Jpn Acad Ser B Phys Biol Sci*. 2009; 85(8): 314-323.
21. Meulmeester E, Ten Dijke P. The dynamic roles of TGF-beta in cancer. *J Pathol*. 2011; 223(2): 205-218.
22. Pfaffl MW, Horgan GW, Dempfle L. Relative expression software tool (REST©) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Res*. 2002; 30(9): e36.
23. Nishikawa S, Ishii H, Haraguchi N, Kano Y, Fukusumi T, Ohta K, et al. microRNA-based cancer cell reprogramming technology. *Exp Ther Med*. 2012; 4(1): 8-14.
24. Miyoshi N, Ishii H, Nagai K, Hoshino H, Mimori K, Tanaka F, et al. Defined factors induce reprogramming of gastrointestinal cancer cells. *Proc Natl Acad Sci USA*. 2010; 107(1): 40-45.
25. Maherali N, Hochedlinger K. Tgfbeta signal inhibition cooperates in the induction of iPSCs and replaces Sox2 and cMyc. *Curr Biol*. 2009; 19(20): 1718-1723.
26. Lin T, Ambasadhan R, Yuan X, Li W, Hilcove S, Abujarour R, et al. A chemical platform for improved induction of human iPSCs. *Nat Methods*. 2009; 6(11): 805-808.
27. Liao B, Bao X, Liu L, Feng S, Zovoilis A, Liu W, et al. MicroRNA cluster 302-367 enhances somatic cell reprogramming by accelerating a mesenchymal-to-epithelial transition. *J Biol Chem*. 2011; 286(19): 17359-17364.
28. Subramanyam D, Lamouille S, Judson RL, Liu JY, Bucay N, Derynck R, et al. Multiple targets of miR-302 and miR-372 promote reprogramming of human fibroblasts to induced pluripotent stem cells. *Nat Biotechnol*. 2011; 29(5): 443-448.
29. Chaudhury A, Howe PH. The tale of transforming growth factor-beta (TGFbeta) signaling: a soigne enigma. *IUBMB Life*. 2009; 61(10): 929-939.
30. Chapnick DA, Warner L, Bernet J, Rao T, Liu X. Partners in crime: the TGFβ and MAPK pathways in cancer progression. *Cell Biosci*. 2011; 1: 42.
31. Li HL, Wei JF, Fan LY, Wang SH, Zhu L, Li TP, et al. miR-302 regulates pluripotency, teratoma formation and differentiation in stem cells via an AKT1/OCT4-dependent manner. *Cell Death Dis*. 2016; 7: e2078.
32. Xue J, Lin X, Chiu WT, Chen YH, Yu G, Liu M, et al. Sustained activation of SMAD3/SMAD4 by FOXM1 promotes TGF-beta-dependent cancer metastasis. *J Clin Invest*. 2014; 124(2): 564-579.
33. Johnson VL, Scott MI, Holt SV, Hussein D, Taylor SS. Bub1 is required for kinetochore localization of BubR1, Cenp-E, Cenp-F and Mad2, and chromosome congression. *J Cell Sci*. 2004; 117(Pt 8): 1577-1589.
34. Nyati S, Schinske-Sebolt K, Pitchiaya S, Chekhovskiy K, Chator A, Chaudhry N, et al. The kinase activity of the Ser/Thr kinase BUB1 promotes TGF-β signaling. *Sci Signal*. 2015; 8(358): ra1.
35. Zhang YE. Non-smad signaling pathways of the TGF-beta family. *Cold Spring Harb Perspect Biol*. 2017; 9(2). pii: a022129.
36. Cargnello M, Roux PP. Activation and function of the MAPKs and their substrates, the MAPK-activated protein kinases. *Microbiol Mol Biol Rev*. 2011; 75(1): 50-83.
37. Meloche S, Pouyssegur J. The ERK1/2 mitogen-activated protein kinase pathway as a master regulator of the G1- to S-phase transition. *Oncogene*. 2007; 26(22): 3227-3239.
38. Cuenda A, Rousseau S. p38 MAP-kinases pathway regulation, function and role in human diseases. *Biochim Biophys Acta*. 2007; 1773(8): 1358-1375.
39. Brancho D, Tanaka N, Jaeschke A, Ventura JJ, Kelkar N, Tanaka Y, et al. Mechanism of p38 MAP kinase activation in vivo. *Genes Dev*. 2003; 17(16): 1969-1978.
40. Ambrosino C, Nebreda AR. Cell cycle regulation by p38 MAP kinases. *Biol Cell*. 2001; 93(1-2): 47-51.