

## Transcript Analysis of Heat Shock Protein 72 in Vitrified 2-Cell Mouse Embryos and Subsequent *In Vitro* Development

Afroz Habibi, Ph.D.<sup>1</sup>, Naser Farrokhi, Ph.D.<sup>2</sup>, Joaquim Fernando Moreira da Silva, Ph.D.<sup>3</sup>, Ahmad Hosseini, Ph.D.<sup>4\*</sup>

1. Department of Anatomical Sciences, International Branch, Shiraz University of Medical Sciences, Kish Island, Iran
2. Department of Agronomy and Plant Breeding, Faculty of Agriculture, Shahrood University of Technology, Shahrood, Iran
3. Animal Reproduction, Department of Agrarian Sciences, University of the Azores, Angra do Heroísmo, Portugal
4. Cellular and Molecular Biology Researcher Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

\*Corresponding Address: P.O.Box: 19395- 4719, Cellular and Molecular Biology Researcher Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran  
Email: Prof\_Hosseini@yahoo.com

Received: 31/Jul/2012, Accepted: 19/Feb/2013

### Abstract

**Objective:** The aim of the study was to compare the effects of two different concentrations of cryoprotectants by cryotopvitrification on survival, developmental capacity and *Heat shock protein 72 (Hsp72)* expression of two-cell mouse embryos.

**Materials and Methods:** In this experimental study, transcript analysis of *Hsp72* gene was performed on non-vitrified and vitrified 2-cell mouse embryos via a nested quantitative polymerase chain reaction (nqPCR) subsequent to normalization with *Hprt1* as the reference gene. The different cryoprotectant combinations were 15% (vit<sub>1</sub>:7.5% of each ethylene glycol (EG) and dimethyl sulfoxide (DMSO)), 30% (vit<sub>2</sub>:15% EG + 15% DMSO) and control group with no cryoprotectants. Vitrified and fresh 2-cell embryos were cultured to obtain cleavage and blastocyst formation rates. The results were analyzed via one-way analysis of variance and the mean values were compared with least significant difference (LSD) ( $p < 0.05$ ).

**Results:** The relative expression of *Hsp72* in vit<sub>2</sub> (30% v/v) was significantly higher than vit<sub>1</sub> (15% v/v). Survival rates were the same for both vitrification treatments and significantly lower than the control group. Cleavage and blastocyst rates in vit<sub>1</sub> were significantly higher than vit<sub>2</sub> while those in two vitrified groups were significantly lower than the control group.

**Conclusion:** Our developmental data demonstrated that vit<sub>1</sub> treatment (7.5% EG and 7.5% DMSO) was more efficient than vit<sub>2</sub> (15% EG and 15% DMSO) in mouse embryos. The cryotopvitrification with two concentrations of cryoprotectants caused the relative changes of *Hsp72* transcript level, but the stability of the gene in vit<sub>1</sub> was significantly higher than vit<sub>2</sub> and closer to the fresh 2-cell embryos.

**Keywords:** Murine, Preimplantation Embryo Development, Quantitative PCR, Vitrification

Cell Journal(Yakhteh), Vol 15, No 4, Winter 2014, Pages: 340-347

**Citation:** Habibi A, Farrokhi N, Moreira da Silva JF, Hosseini A. Transcript analysis of heat shock protein 72 in vitrified 2-cell mouse embryos and subsequent in vitro development. Cell J. 2014; 15(4): 340-347.

### Introduction

In assisted reproduction, embryo cryopreservation has proven to be a powerful tool with applications in bioscience, agriculture and medicine (1). It has been demonstrated that ovarian hyperstimulation syndrome can be decreased via embryo cryopreservation.

Additionally, it has been noted that embryo cryo-

preservation can reduce the occurrence of multiple pregnancies and preserve the fertility of cancer patients (2-4). However, little is known about its molecular impacts on embryos and the future newborn. Hence, considering molecular changes that may occur during and subsequent to embryo cryopreservation would provide a better picture for decision making and managing probable undesirable

outcomes. Evaluating the alterations of particular transcripts that may occur upon cryopreservation or global analysis of transcripts would be the first step towards answering some of the raised questions.

Cryopreservation of embryos usually can be performed either through slow freezing or vitrification. Commonly, a combination of high concentrations of cryoprotectants (typically dimethyl sulfoxide (DMSO) and ethylene glycol (EG) in addition to dehydrating agents such as sucrose or sorbitol have been used in vitrification. Extremely fast cooling of embryos via avoiding ice crystal formation allows vitrification to occur with minimum damage to the cells (4, 5). However, use of high concentrations of cryoprotectants, which are often toxic to the cells may raise some questions regarding the safety issues of this technique (6).

Meanwhile, alterations of vitrification methodology can provide insights to reduce some of its drawbacks. These modifications can be performed either via increasing the cooling rate, a method known as ultra-rapid vitrification (7), or through reducing the vitrification solution volume. It has been suggested that even cells in pure water (without cryoprotectant) can be vitrified if the cooling rate is sufficiently high (8).

Ultra-rapid vitrification methods employ the use of miniature devices, allowing to freeze cells in sub-microlitre volumes (4). Electron microscope grids (9), open-pulled straws (10), cryoloops (11), microdrops (12, 13), cryotops (14), solid surface vitrification (15), nylon mesh (16) and cryotip (17) are amongst successful tools developed in recent years. The approach that minimizes the volume of vitrification solution is the cryotop (14). Cryotop allows loading of very small volume as little as 0.1  $\mu$ l, improving the cooling rate to increase to 23,000  $^{\circ}$ C/minute. Consequently, higher cooling rate allows to useless concentrated solutions and eventually lessening any potentially toxic effects (17, 18).

Expression of many genes including *Heat shock protein (HSP)* family, as its name indicates, is mainly affected in response to the changes in temperature (19). It was previously reported that the expression of *Hsp72/Hsp73* is increased at the 2-cell stage (20). Accordingly, 2-cell mouse embryos were cryopreserved in the presence of two concentrations of cryoprotectants (30 and 15%) and subsequent changes of *Hsp72* and *Hprt1* (house-

keeping gene) were analyzed upon thawing. Cryotop was the instrument of choice for vitrification. Vitrified and fresh 2-cell embryos were cultured to obtain cleavage and blastocyst formation rates. The aim of the study was to compare the effects of two different concentrations of cryoprotectants by cryotop vitrification on survival, developmental capacity and *heat shock protein 72 (Hsp72)* expression of two-cell mouse embryos.

## Materials and Methods

This was an experimental study. This project was approved by the Ethics Committee of Shahid Beheshti University of Medical Sciences in 2009. All chemicals were purchased from Sigma Chemical (St Louis, MO, USA) unless it has been stated otherwise.

CD1 (ICR) female mice aged 8-10 weeks and male mice aged 10-12 weeks (Lisbon University, Portugal) were housed in polycarbonate cages (12 hours light/dark,  $22 \pm 2^{\circ}$ C), and were fed with standard food and fresh water. In all procedures, mice were handled according to the rules stipulated by the Animal Care in Portugal.

### Preparation of 2-cell embryo

Female mice were super ovulated by intraperitoneal injection of 10 IU pregnant mare serum gonadotropin (PMSG), followed by 10 IU of human chorionic gonadotropin (hCG) with a 48 hours interval. Female and male mice (1:1) were mated and checked for vaginal plugs the next morning. The plug-positive female mice were sacrificed by cervical dislocation at 48 hours post-hCG injection (4, 21), and 2-cell embryos were collected by flushing oviducts into potassium simplex optimized medium (KSOM<sup>+</sup>) (Millipore, MA, USA) supplemented with 4 mg/ml bovine serum albumin (BSA) and 20 mM N-2-Hydroxyethylpiperazine-N'-2-Ethanesulfonic Acid (Hepes) buffer (5, 22).

### Study groups

The embryos were vitrified in two different concentrations of cryoprotectants by Cryotop and the changes of *Hsp72* expression, survival, cleavage and blastocyst formation rates in vitrified and non-vitrified groups were assessed. The embryos from the mice sacrificed on each day were collected and then divided into two main groups, vitrified

and control (non-vitrified) groups: the vitrified group was divided into two subgroups vit<sub>1</sub> (15% v/v: 7.5% DMSO+7.5% EG) and vit<sub>2</sub> (30% v/v: 15% DMSO+15% EG). Finally, 195 embryos of vitrified and control groups were evaluated for survival, cleavage and blastocyst rates. 200 embryos were assessed for expression of *Hsp72* and *Hprt1* as the reference gene (23, 24).

For gene expression, each embryo pool containing 10 embryos was stored at -80°C in a minimum volume (2 µl) of RNase free water (23). Experiments in each series were repeated at least three times.

### ***Vitrification and thawing solutions***

As the basal medium or washing solution (WS), modified Dulbecco's phosphate-buffered saline solution containing 10% (v/v) fetal bovine serum (GIBCO, CA, USA) was used. The equilibration solution contained 7.5% (v/v) EG and 7.5% (v/v) DMSO in basal medium.

There were two vitrification solutions (VS) for two vitrified groups, VS<sub>1</sub>: 7.5% (v/v) EG, 7.5% (v/v) DMSO and 0.5 mol/l sucrose in basal medium and VS<sub>2</sub>: 15% (v/v) EG, 15% (v/v) DMSO and 0.5 mol/l sucrose in basal medium. Thawing solution contained 0.5 M sucrose and diluent solutions (D1, D2, D3, D4, and D5) contained 0.4, 0.3, 0.2, 0.1 and 0.05 M sucrose in basal medium, respectively.

### ***Vitrification and thawing***

Two concentrations of vitrification solutions were used to vitrify the mouse 2-cell embryos using Cryotop. Embryos of vit<sub>1</sub> and vit<sub>2</sub> groups were placed in three droplets of equilibration solution for 1 minute total for all of the drops at 25°C. Subsequently, embryos were transferred into vitrification solution VS1 and VS2 respectively for less than 30 seconds. Embryos (6) were moved on the Cryotop (<1 µl vitrification solution) and the Cryotop was immediately submerged in filter-sterilized liquid nitrogen and kept for at least 7 days.

Samples were thawed by plunging the Cryotop into 1 ml of thawing solution at 37°C for 1 minute. Rehydration and gradual removal of cryoprotectants were performed in D1, D2, D3, D4 and D5 for 3 minutes at every step. Thawed embryos were then washed three times in basal medium (Dul-

becco's phosphate-buffered saline solution) for 5 minutes at 25°C.

### ***Definition of morphological survival***

Embryos were defined "morphologically survived", if the embryos possessed an intact zona pellucida, blastomeres and refractive cytoplasm (25, 26). Following the thawing and cryoprotectant removal steps, embryos in 100 µl of sterilized KSOM+AA (Millipore, MA, USA) supplemented with 4 mg/ml BSA were incubated under mineral oil with the availability of 5% (v/v) CO<sub>2</sub>, 5% (v/v) O<sub>2</sub>, and 90% (v/v) N<sub>2</sub> for 1 hour at 37°C.

The validity of morphological classification was confirmed by vital staining with 0.4% sterilized trypan blue solution, a plasma membrane specific dye, in Hanks' balanced salt solution (HBSS) (27, 28). The embryos were examined under an inverted micromanipulation microscope (Eppendorf, NY, USA). The dead cells were stained dark blue by trypan blue but viable cells were able to repel the dye and were not stained. They were counted and recorded as survival rates. Visually dead embryos were discarded, and the morphologically intact embryos were cultured and the gene expression pattern was analyzed.

### ***Embryo culture***

The survived embryos in control, vit<sub>1</sub> and vit<sub>2</sub> groups were cultured in 20 µl droplets of KSOM<sup>+AA</sup> supplemented with 4 mg/ml BSA under mineral oil at 37°C in an atmosphere of 5% CO<sub>2</sub>, 5% O<sub>2</sub> and 90% N<sub>2</sub> to develop into blastocysts. Embryos were assessed daily to record cleavage and blastocyst formation rates for 4 days.

### ***Gene expression***

The relative quantification of gene transcripts was carried out by real-time PCR. Super Script™ III Platinum® Cells Direct Two-Step Quantitative reverse transcriptase PCR (qRT-PCR) Kit with SYBR® Green (Invitrogen, CA, USA) was used to carry out cDNA synthesis and PCR.

### ***Reverse transcription reaction***

Embryos were lysed in 1 µl lysis enhancer and 10 µl resuspension buffer for every PCR tube, which were incubated at 75°C for 10 minutes in a Thermal Cycler (Applied Biosystems 9700, CA,

USA). To degrade any contaminating DNA, the cell lysates were treated with 5  $\mu$ l DNase I and 1.6  $\mu$ l DNase I buffer (10 $\times$ ) at 25°C for 5 minutes. The embryos were treated with 4  $\mu$ l of 25-mM EDTA and incubated at 70°C for 10 minutes. For first-Strand cDNA Synthesis, 20  $\mu$ l 2 $\times$  RT Reaction Mix and 2  $\mu$ l RT Enzyme Mix were added to each tube which was then incubated at 25, 50 and 85°C for 10, 20 and 5 minutes, respectively.

#### *Nested quantitative polymerase chain reaction*

Sometimes the expressions of some genes are very low, which makes the absolute quantification near to impossible. In such cases a prior polymerase chain reaction (PCR) amplification is required to boost the template level for the following

quantification via Real-Time PCR, a technique called "nested quantitative PCR" or nqPCR for short (29, 30). It is noteworthy to mention that the use of PCR amplicons instead of cDNA for the absolute quantification is not as accurate. However in places where the relative quantification serves the purpose, nqPCR provides enough accuracy. Additionally, considering the number of cells or the quantity of RNA that is used for cDNA synthesis, the expression level can be calculated.

The Primer pairs for each gene were designed, synthesized and validated by Molecular Diagnostic Companies (MDC, Burgess Hill, UK). The primer sequences, annealing temperatures and Gen Bank accession numbers are provided in table 1.

*Table 1: Primers and conditions used for quantification of gene expression by real-time PCR*

Amplicon size(bp)	GenBank accession	Sense primer (5'-3')	Anti-sense primer (5'-3')	T <sub>m</sub> (°C)	Amplicon size (bp)
<i>Hsp72</i>	NM_010479	5'ACGGCATCTTCGAGGTGAA 3'	5' TGGTCTGGCTGATGTCCTTCT 3'	50	129
<i>Hprt1</i>	NM_013556	5'TCCTCCTCAGACCGCTTTT3'	5'AGGTATACAAAACAAATCTAGGTCAT3'	48	118

Real-time PCR was conducted in a real-time cycler (Applied Biosystems 7500, CA, USA). To confirm the specificity and integrity of the PCR products, melting curve analyses were performed for all real-time PCR reactions. Standard curves were generated using serial dilutions of cDNAs. The cDNA of each sample was used as template for the preliminary PCR by AmpliTaq Gold PCR Master Mix according to the manufacturer's instruction. Reactions were performed in a final volume of 50  $\mu$ l. The first-round PCR mix contained 2  $\mu$ l specific primer mix (300 nM), 25  $\mu$ l master mix, 5  $\mu$ l cDNA and 18  $\mu$ l sterile water.

The first-round PCR was performed in a thermal cycler (Applied Biosystem 2720, California and USA), by incubation at 95°C for 5 minutes, followed by 30 cycles of 95°C for 15 seconds, specific T<sub>m</sub> for every gene for 15 seconds (Table 1), and 72°C for 60 seconds, and a final extension at 72°C for 7 minutes. The PCR products were separated on 3% agarose gel (pure Nusieve GTC Agarose,

Rockland, USA).

Real time PCR was conducted for cDNA and standards in triplicates with two no-template controls (NTC). Reactions (25  $\mu$ l) contained 12.5  $\mu$ l Platinum<sup>®</sup> SYBR<sup>®</sup> green qPCR super mix-UDG, 0.5  $\mu$ l Rox Reference dye, 0.5  $\mu$ l primer mix (sense and antisense primers, 300 nM each), 6.5  $\mu$ l autoclaved distilled water and 5  $\mu$ l of cDNA in every well.

Cycling parameters were 50°C for 2 minutes (UDG incubation), 95°C for 2 minutes, followed by 50 cycles of 95°C for 15 seconds and 60°C for 30 seconds. The melting curve was analyzed at 95°C for 15 sand temperature lowered to 60°C for 15 seconds. Every experiment was repeated three times.

The data were analyzed with the integrated ABI 7500-V2.0.1 software (Applied Biosystem, California, USA) and were normalized with *Hprt1* within the log linear phase of the amplification curve using the comparative Ct

method (cycle threshold). The relative expression ratio (R) of *Hsp72* was estimated based on a  $\Delta\text{Ct}$  formula (31-33). PCR efficiencies (32, 33) of the genes ranged between 1.98-2.0.  $\Delta\text{Ct}$  was the difference between the Ct values of controls and samples.

### Statistical analysis

One-way analysis of variance (ANOVA) was performed on the average percentages of survived, cleaved embryos, blastocyst formation and relative amount of *Hsp72* mRNA in control, vit<sub>1</sub> and vit<sub>2</sub> groups. Following the analysis of variance, mean values were compared. The

level of significance was set at less than 0.05 and least significant difference (LSD) test was used to compare treatments.

## Results

### Developmental competence of 2-cell embryos following vitrification

In total, 195 in-vivo embryos at 2-cell stage were evaluated for survival, cleavage and blastocyst rates in control, vit<sub>1</sub> and vit<sub>2</sub> groups. The survival rates of vitrified and control groups are summarized in table 2, with no difference between vitrified groups and significantly lower than control ( $p < 0.05$ ).

Table 2: The survival rates of 2-cell embryos in control and vitrified groups

Groups	Concentration of cryoprotectans	No. of total embryos	No. of survived embryos	Mean of survival rate (%)	Standard deviation
control	0%	76	73	95.8a	0.06
vit <sub>1</sub>	15% (7.5% EG+7.5% DMSO)	55	41	75.3b	0.13
vit <sub>2</sub>	30% (15% EG+15% DMSO)	64	45	68.6b	0.07

Control; Non-vitrified group, DMSO; Dimethyl sulfoxide and EG; Ethylene glycol, a and b indicate significant difference between control with vitrified groups ( $p < 0.01$ ). Every experiment was repeated three times.

The cleavage rates of embryos (2-cell to morula) in all groups are shown in figure 1. The cleavage rate in control ( $67.1\% \pm 1.6$ ) was significantly higher than vit<sub>1</sub> ( $48.8\% \pm 0.9$ ). Furthermore, the cleavage rate in vit<sub>1</sub> was significantly higher than vit<sub>2</sub> ( $36.8\% \pm 1.2$ ) groups ( $p < 0.05$ ).




Fig 1: Mean of cleavage rates of 2-cell embryos (to morula) in three groups, cont; control (non-vitrified) group, vit<sub>1</sub>; vitrification with 7.5% DMSO and 7.5% EG, vit<sub>2</sub>; vitrification with 15% DMSO and 15% EG. a, b and c indicate the significant differences among control, vit<sub>1</sub> and vit<sub>2</sub> ( $p < 0.01$ ).

The percentages of blastocyst formation in the control, vit<sub>1</sub> and vit<sub>2</sub> groups were  $43.8 \pm 1.4$ ,  $31.7 \pm 0.9$  and  $21.1 \pm 0.8$ , respectively (Fig 2). The differences among the means of control, vit<sub>1</sub> and vit<sub>2</sub> were significant ( $p < 0.05$ ).





Fig 2: The percentages of blastocyst formation of 2-cell embryos in three groups, cont; control (non-vitrified) group, vit<sub>1</sub>; vitrification with 7.5% DMSO and 7.5% EG, vit<sub>2</sub>; vitrification with 15% DMSO and 15% EG. a, b and c indicate the significant differences among control, vit<sub>1</sub> and vit<sub>2</sub> ( $p < 0.05$ ).


### Expression of Hsp72 mRNA

The effect of different concentrations of cryoprotectants on the expression of *Hsp72* in 2-cell embryos was analyzed with nqPCR and the data were normalized against *Hprt1*. *Hsp72* was significantly up-regulated, 12.9 fold in vit<sub>1</sub> and 32.4 fold in vit<sub>2</sub>, when compared to the control group ( $p < 0.05$ , Fig 3). Moreover, the normalized relative expression ratio of *Hsp72* in vit<sub>2</sub> was significantly higher than vit<sub>1</sub> ( $p < 0.05$ ).

Mean inverse Ct values of *Hprt1* had no significant differences between vitrified and control groups ( $p > 0.05$ , Fig 4).



**Fig 3:** The relative quantification of *Hsp72* after normalization by *Hprt1* in 2-cell embryo groups, cont; control (non-vitrified) group, vit<sub>1</sub>; vitrification with 7.5% DMSO and 7.5% EG, vit<sub>2</sub>; vitrification with 15% DMSO and 15% EG. a, b and c indicate the significant differences among control, vit<sub>1</sub> and vit<sub>2</sub> ( $p < 0.05$ ).



**Fig 4:** Mean inverse Ct values of *Hprt1* as the relevant abundance of transcript 2-cell embryo groups, Ct; threshold cycle, cont; control (non-vitrified) group, vit<sub>1</sub>; vitrification with 7.5% DMSO and 7.5% EG, vit<sub>2</sub>; vitrification with 15% DMSO and 15% EG. Bars are indicative of having no significant difference.

### Discussion

Mouse embryos can be cryopreserved efficiently at a wide range of developmental stages, 2-cell, 8-cell, or the morula stage (25, 34). During the previous years, the success rates of vitrification have

been improved by speeding up the cooling rates of cells via minimizing the sample size (vitrification solution and embryos). This increase effectively prohibits the ice crystal formation (1, 25, 35). Despite the fact that vitrification has proved to be useful in many aspects of cryobiology and fertility restoration, possible molecular consequences of vitrification are yet to be addressed properly. Initially, this can be ascertained through detailed molecular studies of genes that are directly involved in response to temperature change and stress response (36, 37).

Association of *Hsp70*, *Hsp27* and *Hsp90* sub-families have been demonstrated in the protection against apoptosis induced by a variety of stimuli such as heat shock, reactive oxygen species and cytoskeletal perturbation (38-40). Amongst the family of *Hsp*, *Hsp72* is reported to be expressed at 2-cell embryos. For this reason, *Hsp72* was considered as representative of the genes that maybe affected during vitrification with a variety of cryoprotectant concentrations. A concentration of a cryoprotectant is considered suitable when the expression pattern and morphological features of the fresh 2-cell embryos can be replicated as close as possible. Indeed, this means that the cryoprotectant has had minimal effects on the cells.

Here, the previously proposed concentration of cryoprotectants (15% DMSO + 15% EG) was compared with the reduced concentration (7.5% DMSO + 7.5% EG) in cryotop vitrification method. Ultimately, their effects on survival and developmental rates and on the expression of *Hsp72* were compared with the control group (non-vitrification).

The results of the present study demonstrated that the survival rates were the same for both vitrification treatments, but the cleavage and blastocyst formation rates in vit<sub>1</sub> (our proposed concentration) were significantly higher than vit<sub>2</sub> for 2-cell mouse embryo. This may suggest reduced vitrification solution toxicity for vit<sub>1</sub> as opposed to vit<sub>2</sub>. Moreover, the survival and development rates of vitrified embryos were significantly lower than non-vitrified embryos. This might be due to the vitrification-thawing treatment of the embryos at an early stage of development and further be the result of poorly developed stress response mechanisms. In con-

trast to our results, vitrification of human oocytes and embryos had no negative effect on survival and developmental rates (14, 17, 18). These dissimilar outcomes can be explained by the differences that are present between mice and human embryonic cells such as size and shape of the cells and membrane permeability (5).

Two other genes that were previously reported to be expressed in 2-cell embryos (41, 42) were also considered for transcript analysis, Gjal (Connexin 43), a gap junction gene (43-45), and Ped genes, a gene family regulating the rate of preimplantation embryonic development and subsequent embryo survival (46-48). However, our attempts to detect any expression of these genes at this stage failed (data not shown). Transcript analysis of *Hsp72* showed an up-regulation in vitrified groups when compared to the control group, similar to the previous results following other vitrification methods (25, 49). Furthermore, the relative quantification of *Hsp72* in vit<sub>1</sub> was significantly lower than vit<sub>2</sub> and closer to the fresh 2-cell embryos. The fact that *Hsps* play a protective role during imposed stresses to the cells, suppressing several forms of cell death, including apoptosis (50) may suggest that vit<sub>1</sub> treatment had a lesser impact on the overall well-being of the cell. In general, it can be said that 2-cell mouse embryos have experienced thermal stress during vitrification steps, but the concentrated cryoprotectants causes a pronounced stress to the embryos.

## Conclusion

Our developmental data show that cryotopvitrification with 7.5% EG and 7.5% DMSO was more efficient than that with 15% EG and 15% DMSO. Although vit<sub>1</sub> treatment had lower survival and developmental rates compared to the control group, it demonstrated better stability compared with vit<sub>2</sub> based on the *Hsp72* transcript analysis, supporting developmental data.

## Acknowledgements

This study was financially supported by CITA-A, University of the Azores, Angra do Heroísmo, Portugal and Cellular and Molecular Biology Research Center (CMBRC), Shahid Beheshti Uni-

versity of Medical Sciences and Health Services, Tehran, Iran. There is no conflict of interest in this article.

## References

1. Kasai M. Advances in the cryopreservation of mammalian oocytes and embryos: development of ultrarapidvitrification. *Reprod Med Biol.* 2002; 1(1): 1-9.
2. Kawamura T, Motoyama H, Yanaihara A, Yorimitsu T, Arichi A, Karasawa Y, et al. Clinical outcomes of two different endometrial preparation methods for cryopreserved-thawed embryo transfer in patients with a normal menstrual cycle. *Reprod Med Biol.* 2007; 6(1): 53-57.
3. Orief Y, Dafopoulos K, Schultze-Mosgau A, Al-Hasani S. Vitrification: will it replace the conventional gamete cryopreservation techniques?. *Middle East Fertil Soc J.* 2005; 10(3): 171-184.
4. Sheehan CB, Lane M, Gardner DK. The CryoLoop facilitates re-vitrification of embryos at four successive stages of development without impairing embryo growth. *Hum Reprod.* 2006; 21(11): 2978-2984.
5. Moore K, Bonilla AQ. Cryopreservation of mammalian embryos: the state of the art. *Annu Rev Biomed Sci.* 2006; 8: 19-32.
6. Mukaida T, Wada S, Takahashi K, Pedro PB, An TZ, Kasai M. Vitrification of human embryos based on the assessment of suitable conditions for 8-cell mouse embryos. *Hum Reprod.* 1998; 13(10): 2874-2879.
7. Tan J. Vitrification of human oocytes and bovine oocytes and embryos. Ph.D. Westmount, Quebec. Selwyn House School. 2004.
8. Almasi-Turk S, Roozbehi A, Aliabadi E, Haeri A, Sadeghi Y, Hosseini A. Developmental consequences of mouse cryotop-vitrified oocyte and embryo using low concentrated cryoprotectants. *Iran J Reprod Med.* 2009; 7(4): 181-188.
9. Martino A, Songsasen N, Leibo SP. Development into blastocysts of bovine oocytes cryopreserved by ultra-rapid cooling. *Biol Reprod.* 1996; 54(5): 1059-1069.
10. Vajta G, Booth PJ, Holm P, Greve T, Callesen H. Successful vitrification of early stage bovine in vitro produced embryos with the open pulled straw (OPS) method. *Cryo-Lett.* 1997; 18(3): 191-195.
11. Lane M, Schoolcraft WB, Gardner DK. Vitrification of mouse and human blastocysts using a novel cryoloop container-less technique. *Fertil Steril.* 1999; 72(6): 1073-1078.
12. Landa V, Teplá O. Cryopreservation of mouse 8-cell embryos in microdrops. *Folia Biol (Praha).* 1990; 36(3-4): 153-158.
13. Papis K, Shimizu M, Izaike Y. A highly efficient modified vitrification method, for day 3 in vitro produced bovine embryos. *Cryo-Lett.* 1999; 20(4): 203-206.
14. Kuwayama M, Kato O. All round vitrification for human oocytes and embryos. *J Assist Reprod Genet.* 2000; 17: 477-485.
15. Dinnyés A, Dai Y, Jiang S, Yang X. High developmental rates of vitrified bovine oocytes following parthenogenetic activation, in vitro fertilization, and somatic cell nuclear transfer. *Biol Reprod.* 2000; 63(2): 513-518.
16. Matsumoto H, Jiang JY, Tanaka T, Sasada H, Sato E. Vitrification of large quantities of immature bovine oocytes using nylon mesh. *Cryobiology.* 2001; 42(2): 139-144.
17. Kuwayama M, Vajta G, Ieda S, Kato O. Comparison of open and closed methods for vitrification of human embryos and the elimination of potential contamination. *Reprod*

- Biomed Online. 2005; 11(5): 608-614.
18. Kuwayama M. Highly efficient vitrification for cryopreservation of human oocytes and embryos: the Cryotop method. *Theriogenology*. 2007; 67(1): 73-80.
  19. Sonna LA, Fujita J, Gaffin SL, Lilly CM. Effects of heat and cold stress on mammalian gene expression. *J Appl Physiol*. 2002; 92(4): 1725-1742.
  20. Liu CH, Yang CC, Lin DP, Wu MH, Tsai KJ. Stored of Hsp72/Hsp73 in germinal vesicle-stage mouse oocytes. *Reprod Domest Anim*. 2004; 39(1): 19-24.
  21. Uechi H, Tsutsumi O, Morita Y, Takai Y, Taketani Y. Comparison of the effects of controlled-rate cryopreservation and vitrification on 2-cell mouse embryos and their subsequent development. *Hum Reprod*. 1999; 14(11): 2827-2832.
  22. Mukaida T, Takahashi K. Vitrification of day 2-3 human embryos: using various techniques (Cryoloop, Cryotop, and conventional cryostraw). In: Tucker M, Liebermann J, editors. *Vitrification in assisted reproduction*. 1<sup>st</sup> ed. London: Informa healthcare; 2007; 183-193.
  23. Mamo S, Gal AB, Bodo S, Dinnyes A. Quantitative evaluation and selection of reference genes in mouse oocytes and embryos cultured in vivo and in vitro. *BMC Dev Biol*. 2007; 7: 14.
  24. Mamo S, Gal AB, Polgar Z, Dinnyes A. Expression profiles of the pluripotency marker gene POU5F1 and validation of reference genes in rabbit oocytes and preimplantation stage embryos. *BMC Mol Biol*. 2008; 9: 67.
  25. Boonkusol D, Gal AB, Bodo S, Gorhony B, Kitiyanant Y, Dinnyes A. Gene expression profiles and in vitro development following vitrification of pronuclear and 8-cell stage mouse embryos. *Mol Reprod Dev*. 2006; 73(6): 700-708.
  26. Azadbakht M, Valojerdi MR. Development of vitrified-warmed mouse embryos co-cultured with polarized or non-polarized uterine epithelial cells using sequential culture media. *J Assist Reprod Genet*. 2008; 25(6): 251-261.
  27. Briese V. Protocol online. Available from: [http://www-ufk.med.uni-rostock.de/lablinks/protocols/e\\_protocols](http://www-ufk.med.uni-rostock.de/lablinks/protocols/e_protocols). Universitäts-Frauenklinik Rostock. 2002. (28 Oct 2002).
  28. Orynbayeva Z, Kolusheva S, Livneh E, Lichtenshtein A, Nathan I, Jelinek R. Visualization of membrane processes in living cells by surface-attached chromatic polymer patches. *Angew Chem Int Ed Engl*. 2005; 44(7): 1092-1096.
  29. Forsman A, Uzameckis D, Rönnblom L, Baecklund E, Aleskog A, Bindra A, et al. Single-tube nested quantitative PCR: a rational and sensitive technique for detection of retroviral DNA. Application to RERV-H/HRV-5 and confirmation of its rabbit origin. *J Virol Methods*. 2003; 111(1): 1-11.
  30. Munoz-Fernandez MA, Gomez-Chacon GF. Method of in-vitro detection and quantification of HIV DNA by quantitative PCR. In: Patent document. USA: Ladas & Parry LLP; 2008.
  31. Avci ME, Konu O, Yagci T. Quantification of SLIT-ROBO transcripts in hepatocellular carcinoma reveals two groups of genes with coordinate expression. *BMC Cancer*. 2008; 8: 392-402.
  32. Luciano P, Berteza CM, Temporale G, Maffei ME. DNA internal standard for the quantitative determination of hallucinogenic plants in plant mixtures. *Forensic Sci Int Genet*. 2007; 1(3-4): 262-266.
  33. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res*. 2001; 29(9): e45.
  34. Rall WF, Fahy GM. Ice-free cryopreservation of mouse embryos at -196 degrees C by vitrification. *Nature*. 1985; 313(6003): 573-575.
  35. Hiraoka K, Hiraoka K, Kinutani M, Kinutani K. Blastocoele collapse by micropipetting prior to vitrification gives excellent survival and pregnancy outcomes for human day 5 and 6 expanded blastocysts. *Hum Reprod*. 2004; 19(12): 2884-2888.
  36. Beere HM. Stressed to death: regulation of apoptotic signaling pathways by the heat shock proteins. *Sci STKE*. 2001; (93): re1.
  37. Beere HM. The stress of dying: the role of heat shock proteins in the regulation of apoptosis. *J Cell Sci*. 2004; 117(Pt 13): 2641-2651.
  38. Mosser DD, Caron AW, Bourget L, Denis-Larose C, Massie B. Role of the human heat shock protein hsp70 in protection against stress-induced apoptosis. *Mol Cell Biol*. 1997; 17(9): 5317-5327.
  39. Mosser DD, Caron AW, Bourget L, Meriin AB, Sherman MY, Morimoto RI, Massie B. The chaperone function of hsp70 is required for protection against stress-induced apoptosis. *Mol Cell Biol*. 2000; 20(19): 7146-7159.
  40. Beere HM. Death versus survival: functional interaction between the apoptotic and stress-inducible heat shock protein pathways. *J Clin Invest*. 2005; 115(10): 2633-2639.
  41. Houghton FD. Role of gap junctions during early embryo development. *Reproduction*. 2005; 129(2): 129-135.
  42. Dehghani H. Molecular basis of differential gene expression in the mouse preimplantation embryo. *Iran J biotech*. 2007; 5(2): 67-78.
  43. Ruangvoravat CP, Lo CW. Connexin 43 expression in the mouse embryo: localization of transcripts within developmentally significant domains. *Dev Dyn*. 1992; 194(4): 261-281.
  44. Juneja SC, Barr KJ, Enders GC, Kidder GM. Defects in the germ line and gonads of mice lacking connexin43. *Biol Reprod*. 1999; 60(5): 1263-1270.
  45. Dahl E, Winterhager E, Reuss B, Traub O, Butterweck A, Willecke K. Expression of the gap junction proteins connexin31 and connexin43 correlates with communication compartments in extraembryonic tissues and in the gastrulating mouse embryo, respectively. *J Cell Sci*. 1996; 109 (Pt 1): 191-197.
  46. Cai W, Cao W, Wu L, Exley GE, Waneck GL, Karger BL, et al. Sequence and transcription of Qa-2-encoding genes in mouse lymphocytes and blastocysts. *Immunogenetics*. 1996; 45(2): 97-107.
  47. McElhinny AS, Kadow N, Warner CM. The expression pattern of the Qa-2 antigen in mouse preimplantation embryos and its correlation with the Ped gene phenotype. *Mol Hum Reprod*. 1998; 4(10): 966-971.
  48. Wu L, Feng H, Warner CM. Identification of two major histocompatibility complex class Ib genes, Q7 and Q9, as the Ped gene in the mouse. *Biol Reprod*. 1999; 60(5): 1114-1119.
  49. Khodabandeh Z, Amidi F, Nori MH, Sobhani A, Mehrannia K, Abbasi M, et al. Expression of heat shock protein (HSP A1A) and MnSOD genes following vitrification of mouse MII oocytes with cryotopmethod. *Yakhteh*. 2010; 12(1): 113-119.
  50. Neuer A, Mele C, Liu HC, Rosenwaks Z, Witkin SS. Monoclonal antibodies to mammalian heat shock proteins impair mouse embryo development in vitro. *Hum Reprod*. 1998; 13(4): 987-990.