

The Effect of Nano-Titanium Dioxide on Limb Bud Development of NMRI Mouse Embryo *In Vivo*

Kazem Parivar, Ph.D.¹, Nasim Hayati Rudbari, Ph.D.¹, Ramazan Khanbabaee, Ph.D.², Mahya Khaleghi, M.Sc.^{2*}

1. Department of Biology, Faculty of Science, Islamic Azad University, Science and Research Branch, Tehran, Iran

2. Department of Biology, Faculty of Science, Islamic Azad University, Ghaemshahr, Mazandaran, Iran

*Corresponding Address: P.O.Box: 163, Department of Biology, Faculty of Science, Islamic Azad University, Ghaemshahr, Mazandaran, Iran
Email: mahya.khaleghi@gmail.com

Received: 22/Jan/2014, Accepted: 11/Agu/2014

Abstract

Objective: There is a wide application of titanium dioxide (TiO₂) nanoparticles (NPs) in industry. These particles are used in various products, and they also has biological effects on cells and organs through direct contact.

Materials and Methods: In this experimental research, the effect of TiO₂ on chondrogenesis of forelimb buds of mice embryos was assessed in *in vivo* condition. Concentrations of 30, 150 and 500 mg/kg body weight (BW) TiO₂ NPs (20 nm size) dissolved in distilled water were injected intraperitoneally to Naval Medical Research Institute (NMRI) mice on day 11.5 of gestation. On day 15, limb buds were amputated from the embryos and skel-etogenesis of limb buds were studied.

Results: TiO₂ NPs caused the significant changes in chondrocytes in the following developmental stages: resting, proliferating, hypertrophy, degenerating, perichondrium and mesenchymal cells. Decreased number of mesenchymal cells and increased level of chondrocytes were observed after the injection of different concentrations of TiO₂, which proves the unpredictable effects of TiO₂ on limb buds.

Conclusion: Results of the present study showed TiO₂ NPs accelerated the chondrogenesis of limb buds, but further studies are recommended to predict TiO₂ toxicity effects on organogenesis.

Keywords: Titanium Dioxide, Nanoparticles, Limb Bud, Chondrogenesis

Cell Journal(Yakhteh), Vol 17, No 2, Summer 2015, Pages: 296-303

Citation: Parivar K, Hayati Rudbari N, Khanbabaee R, Khaleghi M. The effect of nano-titanium dioxide on limb bud development of NMRI mouse embryo *in vivo*. Cell J. 2015; 17(2): 296-303.

Introduction

Titanium dioxide (TiO₂) is widely used as an industrial nanomaterial. The use of TiO₂ nanoparticles (NPs) as an additives in pharmaceutical and food industries as well as in various products, including coatings, ceramics, anti-fouling paints, cosmetics and sunscreens, has gained increasing attention in past few years (1, 2). The unique physicochemical characteristics of NPs, such as high reactivity, small size, and large surface area per mass, have raised great concerns on the adverse effects of TiO₂ NPs on ecological system and human health (3-5). Due to its widespread use, humans are increasingly exposed to TiO₂ NPs material. The respiratory tract and skin are the main exposure areas. It has been also demonstrated that TiO₂ NPs changes phagocytic activity in cultured macro-phages and the integrity of the cell membrane (6). TiO₂ occurs

primarily in the forms of the minerals rutile, anatase and brookite (7, 8). Therefore, we aimed to evaluate the effects of TiO₂ NPs on chondrogenesis of fore-limbs skeletons of mouse embryo *in vivo*.

Materials and Methods

Animals and treatment

In this experimental research, we studied the effect of TiO₂ on limb bud of mice. This project was carried out at the Animal Biology Laboratory of Islamic Azad University, Science and Research Branch, Tehran, Iran. Naval Medical Research Institute (NMRI) mice (5 males, 20 females, 30 ± 5 g) were purchased from the Pasture Institute, Tehran, Iran. Mice were housed in an animal room at 24 ± 2°C with a 12-hour light/dark cycle for five days, before

starting the experiment. Food and water were provided and copulations were set up. Day one of vaginal plug observation was determined as day 0 of gestation. After 11.5 days, the body weight of animals were weighed and randomly divided into 5 groups (n=3 per group). Three experimental groups were injected intraperitoneally with size 20 nm TiO₂ NPs at concentrations of 30, 150 and 500 mg/kg body weight (BW), a sham group received an injection of 1ml distilled water, and non-injected group was assigned as control group. Four days later, the mice were sacrificed after being anesthetized by ether. All procedures used in animal experiments were in compliance with the Ethics Committee of Science and Research Branch of Islamic Azad University. Embryos were excised from uterus and amniotic membrane, and then both right and left forelimbs were carefully amputated, rinsed with Hank's balanced salt solution (HBSS, Merck, Germany), and prepared for histological and morphological studies.

Histological methods

The tissues were immediately fixed in Bouin's solution (Merck, Germany), for 2 hours. Briefly the samples

were embedded in paraffin blocks, serially sectioned into 5- μ m in thickness and placed onto the glass slides. After hematoxylin-eosin (HE) staining, the slides were observed using an optical microscope (Nikon, USA) (Fig.1). The following measurements were then performed in each groups: length of finger-palm (region 1), length of wrist-forearm (region 2) and arm-forearm (region 3), using scaled graticule. Furthermore number of mesenchymal, perichondrial cells, resting, proliferating, hypertrophic and degenerating chondrocytes cells were counted in 3 mentioned region from 6 fields of microscope views in each groups. The mesenchymal cells contains a large, round nucleus, which is surrounded by long cytoplasm. The perichondrium is a sheath of dense connective tissue that surrounds cartilage in most places. The resting zone consists typical round chondrocytes. In the proliferative zone, oval chondrocytes begin to divide rapidly and form columns of stacked cells parallel to the long axis of the bone. The hypertrophic zone contains swollen chondrocytes appearing after proliferating chondrocytes. The degenerative zone contains disintegrated swollen chondrocytes (Fig.2).




Fig.1: Photomicrograph of limb bud tissues on day 15 of development. Samples of each group were fixed in Bouin's fluid, embedded in paraffin, serially sectioned (5- μ m thick) and stained using hematoxylin-eosin (HE) method. **A.** Control $\times 40$, **B.** Sham $\times 40$, **C.** Concentration of 30, **D.** 150 and **E.** 500 $\times 40$. 1; Mesenchymal cells, 2; Chondrocytes and 3; Finger grooves.




Fig.2: Photomicrograph of limb bud tissues on day 15 of development. **A.** Chondrocytes cells in control sample $\times 100$, **B.** Chondrocytes in sham sample $\times 100$, exposed to distilled water. Chondrocytes in experimental samples $\times 400$, **C.** They were exposed to titanium dioxide (TiO₂) nanoparticles (NPs) at concentrations of 30, **D.** 150 and **E.** 500 mg/kg body weight (BW) on day 11- of development. 1; Resting chondrocytes, 2; Proliferating chondrocytes, 3; Hypertrophic chondrocytes, 4; Degenerating chondrocytes, 5; Perichondrium cells and 6; Mesenchymal cells.

Statistical analysis

The mean values of forelimbs parameters were calculated from six sections per slide in each group (n=6). All data were conducted using the SPSS (SPSS Inc., Chicago, IL, USA) software. One-way ANOVA was used to compare the differences among means belonging to multi-group data. All values were calculated from standard errors of means and statistical significance was considered at level of $P \leq 0.05$.

Results

Morphometric study

In the experimental groups, the limb bud tissue had no abnormal pathological changes compared with the control. Abortion percentage in experimental groups was 8%. Morphological differences were observed in length of forelimb buds and the experimental samples showed a significant decrease in length of finger-palm (region 1), length

of wrist-forearm (region 2), and arm-forearm (region 3), as compared with the control and sham groups (Fig.3).

Histological examinations

Comparison of the forelimb buds sections exposed to different concentrations of TiO₂ NPs confirmed significant changes in cell proliferation and differentiation. Figures 4-9 shows the change of chondrocyte numbers in forelimb buds.

In finger-wrist (region 1), mesenchymal cell counts showed a significant decrease in experimental groups compared with sham and control groups, but perichondrial cells showed a significant increase in comparison with the sham and control groups. Resting chondrocytes counts in this region showed an increase as compared with sham and control groups and this increase was significant in the 30 mg/kg TiO₂ treatment. The number of proliferating chondrocytes showed no significant changes in this region in all treatments.

Hypertrophic chondrocytes showed a significant increase in both 150 and 500 mg/kg TiO₂-treated groups, whereas there were no significant differences in 30 mg/kg TiO₂ treatment in comparison with the sham and control groups. Degenerating chondrocytes were observed in wrist-forearm (region 2) and arm-forearm (region 3).

In wrist-forearm (region 2), mesenchymal cell counts showed a significant decrease in 150 and 500 mg/kg TiO₂-treated groups compared with the sham and control groups, indicating there were no significant differences in 30 mg/kg TiO₂ treatment. Perichondrial cells showed no significant changes in comparison with the sham and control groups, but a significant decrease by exposure to 500 mg/kg TiO₂ were observed. Resting chondrocytes in this region showed a significant decrease compared with the sham and control samples. Proliferating chondrocytes revealed no significant changes compared with the sham and control groups. Hypertrophic chondrocytes counts in all experimental groups showed a significant increase in compared with the sham and control groups. The number of degenerating chondrocytes were reduced by ex-

posure to 500 mg/kg TiO₂, but this decrease was not significant; however, the number of cells by exposure to concentrations of 30 and 150 mg/kg TiO₂ were significantly decreased.

In arm-forearm (region 3), mesenchymal cell counts were significantly higher in 30 mg/kg TiO₂ treatment than exposed to concentrations of 150 and 500 mg/kg TiO₂, showing no significant changes compared with the sham and control groups. Perichondrial cells showed a significant increase in 500 mg/kg TiO₂ treatment in comparison with the sham and control samples. Resting chondrocytes counts showed a significant increases in 30 and 500 mg/kg TiO₂ treatments compared with the sham and control groups. Proliferating chondrocytes showed no significant changes in 30 and 150 mg/kg TiO₂ treatments in comparison with the sham and control groups, but hypertrophic chondrocytes showed no obvious differences in comparison with the sham and control groups. Degenerating chondrocytes were reduced in experimental groups and revealed a decrease with doses of 150 and 500 mg/kg TiO₂, suggesting a significant difference.




Fig.3: Histograms of length of regions in forelimb buds on day 15 of development. Region 1 belonging to finger-palm, region 2 belonging to wrist-forearm and region 3 belonging to arm-forearm. Data are presented as means ± SEM. *, P <0.05, **, P <0.01 and ***, P <0.001.




Fig.4: Histograms of number of mesenchymal cells in control, sham and experimental groups on day 15 of development. Region 1 belonging to finger-palm, region 2 belonging to wrist-forearm and region 3 belonging to arm-forearm region. Data are presented as means \pm SEM. *, P<0.05, **, P<0.01 and ***, P<0.001.




Fig.5: Histograms of number of perichondrium cells in control, sham and experimental groups on day 15 of development. Region 1 belonging to finger-palm, region 2 belonging to wrist-forearm and region 3 belonging to arm-forearm. Data are presented as means \pm SEM. *, P<0.05, **, P<0.01 and ***, P<0.001.




Fig.6: Histograms of number of resting chondrocyte cells in control, sham and experimental groups on day 15 of development. Region 1 belonging to finger-palm region, region 2 belonging to wrist-forearm region and region 3 belonging to arm-forearm region. Data are presented as means \pm SEM. *, P <0.05, **, P<0.01 and ***, P<0.001.




Fig.7: Histograms of number of proliferating chondrocytes in control, sham and experimental groups on day 15 of development. Region 1 belonging to finger-palm region, region 2 belonging to wrist-forearm region and region 3 belonging to arm-forearm region. Data are presented as means \pm SEM. *, $P < 0.05$, **, $P < 0.01$ and ***, $P < 0.001$.




Fig.8: Histograms of number of hypertrophic chondrocytes in control, sham and experimental groups on day 15 of development. Region 1 belonging to finger-palm, region 2 belonging to wrist-forearm and region 3 belonging to arm-forearm. Data are presented as means \pm SEM. *, $P < 0.05$, **, $P < 0.01$ and ***, $P < 0.001$.




Fig.9: Histograms of number of degenerating chondrocytes in control, sham and experimental groups on day 15 of development. Region 1 belonging to finger-palm, region 2 belonging to wrist-forearm and region 3 belonging to arm-forearm. Data are presented as means \pm SEM. *, $P < 0.05$, **, $P < 0.01$ and ***, $P < 0.001$.

Discussion

The smaller size of the NPs causes a greater change in the specific embryonic areas, resulting in stronger reaction activity. These characteristics can reduce the stability of the cell membrane that leads to cell injury. It can also influence the activity and function of cells through interaction with macromolecules (9). There are great attention to toxicity of TiO₂ NPs, but data are limited. Our results showed that the embryonic developmental processes were influenced by intraperitoneal injections of various doses of TiO₂ NPs solutions. There are various studies regarding the potential neuronal uptake, translocation of inhaled particulates and pathological effects on the brain (10, 11). Researchers reported that anatase TiO₂ NPs injected at the mouse abdominal cavity migrated anteriorly into the brain that resulted in the oxidative stress and injury of the brain, and subsequently disturbed the normal metabolism of neurochemicals (12). Our study showed that TiO₂ NPs migrated into the uterus, affected the embryos, and caused the abortion. We observed a significant decrease in length of regions, suggesting that this reduction were dose-dependent. NPs (TiO₂ or gold) are no longer distributed in the cytoplasm after being internalized by cells, so they are preferentially placed in mitochondria. When the mitochondria are invaded by NPs, the antioxidant defense ability may be changed (12). In another study investigated the biochemical parameters after daily injection of 5 nm anatase TiO₂ NPs into the abdominal cavity of imprinting control region (ICR) mice for 14 days. The antioxidative responses observing in liver were reduced and hepatic lipids peroxide were increased in mice when using TiO₂ NPs that suggested an oxidative attack (13). We worked on mice were injected on day 11 of gestation that was the critical point in limb bud development. We selected limb bud due to its variation in development and morphological changes. Results showed that TiO₂ Nps caused different changes in chondrogenesis that led to a decrease in the number of mesenchymal cells that may be due to their differentiation into chondrocytes, suggesting a positive effect of TiO₂ Nps on these cells. Furthermore, our findings revealed significant changes in resting, proliferating, and hypertrophic zones. Therefore TiO₂ NPs accelerated the development of limb buds. Resting chondrocyte counts showed an increase as

compared with the sham and control groups which means Nps prevented the differentiation of these cells in regions 1 and 3. The number of proliferating chondrocyte cells was insignificant in region 1 and 2, whereas surprisingly in region 3 that the mice were treated with 500 mg/kg TiO₂ Nps, the number of cells showed a significant increase, suggesting that TiO₂ Nps increased the proliferating chondrocyte cells by disrupting the cells cycle. So, the exposure dose is an important parameter in inducing toxicity. Genetically *fibroblast growth factors (FGFs)* play a crucial role in early step of chondrogenesis. These factors bind to their tyrosine kinase receptors and regulate proliferation and differentiation. Another tests on mice revealed requirements for *bone morphogenetic protein (BMP)* signaling pathways in multiple aspects of chondrogenesis such as proliferation and differentiation of cells and also demonstrated that progression of chondrocytes is controlled by the balance between signaling outputs from *BMP* and *FGFs* pathways (14, 15). Additionally *parathyroid hormone related protein (PTHrP)* along with *Indian hedghog (Ihh)* regulate cells from proliferation to hypertrophied stages by formation a negative feedback loop (16). Hypertrophic chondrocytes count in regions 1 and 2 showed a significant increase that may be due to differentiation of proliferating chondrocytes to hypertrophic chondrocytes that is in contrast to region 3, showing no obvious differences in comparison with the sham and control groups. Perichondrial cells in TiO₂-treated groups showed a significant increase in comparison with the control group because perichondrium is a type of irregular collagenous connective tissue that plays a role in growth and repair of cartilage. During aggregation of mesenchymal cells by *SOX9* expression, cells in the center were committed to differentiate into cartilage, while cells in the periphery remained undifferentiated and appeared in form of perichondrium status (17). In a study that mice treated with the doses of 125 and 250 mg/kg BW anatase TiO₂ NPs for consecutive 30 days displayed a decrease in body weight, seriously damaged liver function, as well as increased coefficients of the liver, kidney, spleen and thymus. It is very likely that liver function damage in mice is caused by higher anatase TiO₂ NPs that is closely associated with the damage of haemostasis blood system and immune response because dose of 62.5 mg/kg TiO₂ NPs has little influence on haemosta-

sis blood system and immune response in mice (18). Another study showed high-dose of anatase TiO₂ NPs (5 nm) through intraperitoneal injection could damage liver function (19). In our study, mesenchymal cells counts were significantly higher in the 30 mg/kg TiO₂ treatment that led to the increased level of resting chondrocytes in regions 1 and 3, indicating that exposure dose have important role in toxicity. In an experiments rats were intra-tracheally instilled with 0.5, 5, or 50 mg/kg of 5, 21 and 50 nm TiO₂ primary particles and their results showed that 5 and 21 nm TiO₂ can induce pulmonary lesions when exposure dose is >5.0 mg/kg TiO₂ particles. It is noted that if the exposure dose is ≥50 mg/kg, 5 nm TiO₂ primary particles may suppress the phagocytotic ability of alveolar macrophages (AMs). Our results also confirmed the important roles of particle size and exposure dose. Further studies are needed to elucidate the underlying mechanisms and thier correlation with the physico-chemical properties of nano-TiO₂ (20).

Conclusion

TiO₂ Nps could accelerate the development of limb buds in specific dose and particle sizes. We suggest a chondrogenic potential of TiO₂ NPs which may be usefull in genetic abnormalities, although the toxicity of different concentrations of TiO₂ NPs on organogenesis should be investigated further.

Acknowledgments

The authors would like to gratefully thank all those who have helped in this research. There is no conflicts of interest in this study.

References

- Jin CY, Zhu BS, Wang XF, Lu QH. Cytotoxicity of titanium dioxide nanoparticles in mouse fibroblast cells. *Chem Res Toxicol.* 2008; 21(9): 1871-1877.
- Vamanu CI, Cimpan MR, Hol PJ, Sornes S, Lie SA, Gjerdet NR. Induction of cell death by TiO₂ nanoparticles: studies on a human monoblastoid cell line. *Toxicol In Vitro.* 2008; 22(7): 1689-1696.
- Erb U, Aust KT, Palumbo G. Electrodeposited nanocrystalline materials. In: Koch CC, editor. 2nd ed. *Nanostructured materials: processing, properties and potential applications.* New York: Noyes; 2002; 179-222.
- Oberdorster G, Maynard A, Donaldson K, Castranova V, Fitzpatrick J, Ausman K, et al. Principles for characterizing the potential human health effects from exposure to nanomaterials: elements of a screening strategy. *Part Fibre Toxicol.* 2005; 2: 8.
- Warheit DB, Webb TR, Reeda KL, Frerichs S, Sayes CM. Pulmonary toxicity study in rats with three forms of ultrafine-TiO₂ particles: differential responses related to surface properties. *Toxicology.* 2007; 230(1): 90-104.
- Zhang Q, Kusaka Y, Donaldson K. Comparative injurious and proinflammatory effects of three ultrafine metals in macrophages from young and old rats. *Inhal Toxicol.* 2000; 12 Suppl 3: 267-273.
- Wakefield G, Stott J, Hock J. Sunscreens and cosmetics containing manganese doped titanium oxide nanoparticles. *SOFW J.* 2005; 131: 46-51.
- Zhang H, Bandfield JF. Thermodynamic analysis of phase stability of nanocrystalline titania. *J Mater Chem.* 2008; 8: 2073-2076.
- Bermudez E, Mangum JB, Wong BA, Asgharian B, Hext PM, Warheit DB, et al. Pulmonary responses of mice, rats, and hamsters to subchronic inhalation of ultrafine titanium dioxide particles. *Toxicol Sci.* 2004; 77(2): 347-357.
- Oberdorster G, Sharp Z, Atudorei V, Elder A, Gelein R, Kreyling W, et al. Translocation of inhaled ultrafine particles to the brain. *Inhal Toxicol.* 2004; 16(6-7): 437-445.
- Dorman DC, Struve MF, James RA, Marshall MW, Parkinson CU, Wong BA. Influence of particle solubility on the delivery of inhaled manganese to the rat brain: manganese sulfate and manganese tetroxide pharmacokinetics following repeated (14-day) exposure. *Toxicol Appl Pharmacol.* 2004; 170(2): 79-87.
- Ma L, Liu J, Li N, Wang J, Duan Y, Yan J, et al. Oxidative stress in the brain of mice caused by translocated nanoparticulate TiO₂ delivered to the abdominal cavity. *Biomaterials.* 2010; 31(1): 99-105.
- Liu H, Ma L, Liu J, Zhao J, Yan J, Hong F. Toxicity of nano-anatase TiO₂ to mice: liver injury, oxidative stress. *Toxicol Environ Chem.* 2010; 92(1): 175-186.
- Yoon BS, Pogue R, Ovchinnikov DA, Yooshi I, Mishina Y, Behringer RR, et al. BMPs regulate multiple aspects of growth-plate chondrogenesis through opposing actions on FGF pathway. *Development.* 2006; 133(23): 4667-4678.
- Pizette S, Niswander L. Early steps in limb patterning and chondrogenesis. *Novartis Found Symp.* 2001; 232: 23-36.
- Vortkamp A, Lee K, Lanske B, Segre GV, Kronenberg HM, Tabin CJ. Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein. *Science.* 1996; 273(5275): 613-622.
- Dudley A, Ros MA, Tabin CJ. A re-examination of proximodistal patterning during vertebrate limb development. *Nature.* 2002; 418(6897): 539-544.
- Duan Y, Liu J, Ma L, Li N, Liu H, Wang J, et al. Toxicological characteristics of nanoparticulate anatase titanium dioxide in mice. *Biomaterials.* 2010; 31(5): 894-899.
- Liu H, Ma L, Zhao J, Liu J, Yan J, Ruan J, et al. Biochemical toxicity of mice caused by nano-anatase TiO₂ particles. *Biol Trace Elem Res.* 2009; 129(1-3): 170-180.
- Liu R, Yin L, Pu Y, Liang G, Zhang J, Su Y, et al. Pulmonary toxicity induced by three forms of titanium dioxide nanoparticles via intra-tracheal instillation in rats. *Prog Nat Sci.* 2009; 19(5): 573-579.