

An Immunohistochemical Study of Retinal Collagen IV Expression during Pre- and Postnatal Periods in Balb/c Mice

Mehdi Jalali, Ph.D.¹, Mohammad Reza Nikravesh, Ph.D.^{1*}, Abbas Ali Moeen, Ph.D.²,
Mohammad Hassan Karimfar, Ph.D.³, Shahin Saidi Nejat, M.D.¹, Shabnam Mohammadi, M.Sc.¹,
Houshang Rafighdoost, Ph.D.⁴

1. Department of Anatomy, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
2. Department of Anatomy, School of Medicine, Zabol University of Medical Sciences, Zabol, Iran
3. Department Anatomy, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran
4. Department of Anatomy, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

* Corresponding Address: P.O.Box: 9138813944, Department of Anatomy, School of Medicine,
Mashhad University of Medical Sciences, Mashhad, Iran
Email: doctornikravesh@yahoo.com

Received: 24/Jun/2009, Accepted: 2/Jan/2010

Abstract

Objective: Basement membranes are specialized extracellular matrices which play important roles such as cell regulation, proliferation and migration. Collagen fibers, especially type IV, are the most important basement membrane constituents. As retina is one of the target organs in diabetes mellitus, and nephropathy is a major cause of end stage renal and retinal diseases resulting in increased morbidity and mortality, early diagnosis leads to better treatment. Hence, in this investigation, the appearance and distribution of collagen IV during gestational days and early postnatal periods were observed.

Materials and Methods: 24 intact female Balb/c mice were kept under normal conditions. After mating, appearance of a vaginal plug was assumed as day zero of the pregnancy. From days 13-18 of gestation, the pregnant mice were euthanised and their embryos as well as pups from days 1 to 5 were collected. For histochemical studies, heads of the specimens were fixed, serially sectioned and immunohistochemical studies were performed by using monoclonal antibodies for tracing of collagen type IV.

Results: Our findings revealed that the amount of collagen IV in the internal limiting membrane (ILM) and extra cellular matrix (ECM) of the retina, as well as vessels of the vitreous body appear on embryonic day 16. Also, a patchy distribution was observed in the pigmented epithelium which continued to further develop until the end stage of embryonic life. Strong labeling was observed until postnatal day 3 but did not increase significantly thereafter.

Conclusion: These findings establish the importance of collagen IV during the critical period of retinal development. In addition, this study indicates that high levels of collagen IV are present in the basal membrane (BM) of the inner limiting membranes and pigmented epithelium (3rd post natal on the 3rd postnatal day).

Keywords: Collagen IV, Extracellular Matrix, Development, Retina, Mouse

Cell Journal(yakhteh), Vol 12, No 4, Winter 2011, Pages: 447-452

Introduction

The basement membrane (BM) is a specialized region of the extra cellular matrix that consists of different components such as collagen, laminin, fibronectin, sulfated and nonsulfated glycosaminoglycan (1). Among BM components, fibronectin, heparin sulfate and collagen type IV play critical roles in cell signaling and differentiation of the optic cup (2-4). In other words, major components of conducting signals in retinal development are also components of the extra cellular

matrix (ECM) which have been organized into the basement membrane (5, 6). Some of these molecules bind to other cells, whereas others bind to the basement membrane of the developing retina where they are called the limiting membranes (7). Hence, it could be speculated that certain molecules and components of extra cellular matrix are necessary for cell differentiation (8, 9). Among these macromolecules, collagen type IV plays a major regulatory role in cell behaviors and developmental phenomena such as proliferation, migra-

tion, differentiation, morphogenesis and metabolism (10-12). The most prominent roles of ECM are migration and adhesion of cells to cells and their substrates (13). Collagen type IV serves these roles in both inner and outer limiting membranes of the retina and optic cup (14). Also, components of ECM such as the α_1 - α_2 chains of collagen type IV are required for controlling blood circulation in the developing visual system (15) as this molecule is known to have an important role in eye physiology. For example, transparent areas of the eye are dependent on the correct development of ECM and special arrangement of collagen (16). Collagen type IV plays an important role in the morphogenesis and cell association during retinal development. It seems that appearance and synthesis of collagen can represent the development and differentiation of retina (17). In other words, morphogenesis of embryonic tissues such as the retina is made possible by intracellular mechanisms modulated by exogenous signals (18). In fact, growth factors and extra cellular matrix components act as mediators in cell development and differentiation (19). Because collagen type IV is an important component of extracellular and basement membranes (20), the aim of the presented study was to determine its appearance and distribution by labeling and tracing it during the critical period of optic cup and retinal development until the last day of pregnancy as well as after birth to more precisely investigate its role in retinal differentiation.

Materials and Methods

Animal procedure

In this study, 24 female Balb/c mice were obtained from animal house of Mashhad University of Medical Sciences. The environmental conditions were 23-25°C, relative humidity of 50-55% and 12 hour light-dark cycles. Mature females were mated overnight with males of the same strain (two females and one male) and examined the next morning for vaginal plugging. Females with vaginal plugs were designated day zero of pregnancy. Pregnant mice from gestational days 13-19 (E13 to E19) and pups of 1 to 5 postnatal days of age were euthanized and transcardinally perfused with physiological saline followed by formalin (Merck Pharmaceuticals, Germany); their brains were removed and postfixed for at least 24 hours at room temperature in the same fixative. The head of embryos and their removed eyeballs were routinely processed and embedded in paraffin with eight-micron sections being carried out serially. Only slides containing eyeball samples were used in this study. The sections were stained with

cresyl violet or incubated with the anti-collagen IV monoclonal antibody (Sigma-Aldrich, USA). Photographs were taken using an Olympus BH2 light microscope.

Immunohistochemistry study

The avidin-biotin-peroxidase complex (ABC) was used for our immunohistochemical study. Sections were deparaffinized, rehydrated and washed twice for 5 minutes in a 0.05 M Tris buffer containing 1.5% NaCl, pH: 7.4. To block nonspecific antibodies, sections were preincubated in 0.3% Triton X-100 in TB-NaCl followed by 5% goat serum (Gibco, UK) for 2-3 hours; they were then left for 12-24 hours at 4°C to react with primary antibody anti-collagen (Sigma-Aldrich, USA) diluted 1:250 in TB-NaCl with 0.3% Triton X-100 and 2% bovine serum. The tissues were then washed three times with TB-NaCl for 10 minutes each time and incubated for 2 hours in biotinylated goat anti-rabbit IgG (1:400 in TB-NaCl). After three further rinses, each lasting 1 hour, the endogenous peroxidase activity was blocked by incubating the sections in 0.03% H₂O₂ in methanol for 30 minutes. The tissues were then incubated for 2 hours in 1:100 avidin-biotinylated horseradish peroxidase complex, each washed in TB-NaCl for 30 minutes three times, and finally reacted with a 0.03% solution of 3, 3'-diaminobenzidine tetrahydrochloride (Sigma-Aldrich, USA) containing 0.03% H₂O₂ for 10-15 minutes before they were washed and lightly counterstained with hematoxylin and eosin (Sigma-Aldrich, USA). Subsequently, the tissues were washed again, air-dried, dehydrated and then mounted in PBS/glycerol (1:9). Using an Olympus BX51 light microscope, images were taken and the lens collagen IV reactions in different embryonic stages were evaluated.

Statistical analysis

The staining intensity was graded by two different individuals in accordance with Firth's method. The scoring gradation had four stages and reflected the severity of reaction from negative to weak, moderate, strong and very strong. P-values < 0.05 were considered as significant.

Results

The immunohistochemical data collected until the early 15th day of gestation indicated no detectable collagen IV reaction at any area of the eyeball (Fig 1A). The first noticeable change was the appearance of a very pale reaction on the late 15th day of gestation which was distributed in the extra cellular matrix proteins of the retinal ganglion cells and

vitreous body (Fig 1B). On the 16th day of gestation, the retinal inner and external limiting membranes as well as hyaloid vessels of the vitreous body showed weak reactions (Fig 1C, D). Also, a patchy distribution was observed in the retinal pigment epithelium. This reaction continued and slightly increased in the following days (Table1, Fig 2). Strong labeling was observed in the first 3 postnatal days, but after this period, collagen IV reaction did not show distinct changes, further intensity of reaction was observed in the inner limiting membrane and pigmented epithelium.

Table 1: Distribution of collagen IV during retinal development in mouse ED (embryonic day), PND (postnatal day), ILM (inner limiting membrane), ECM (extra cellular matrix) and ELM (external limiting membrane). Collagen IV reaction intensity: no reaction (-), weak (+), moderate (++), high (+++) and severe (++++).

DE	ILM	ECM	ELM	BM/pigment epithelial cells
14	(-)	(-)	(-)	(-)
15	(+)	(-)	(+)	(++)
17	(++)	(+)	(++)	(+++)
3 rd PN	(++++)	(+++)	(++)	(++++)




Fig 1: Retinal cross sections (15th gestation day) incubated with the collagen IV antibody (A) and with higher magnification (B), extra cellular matrix (ECM) and inner limiting membrane (ILM) indicate no reaction (arrows). On day 16 and 18 of gestation (C and D respectively) arrow-marked areas showed weak reaction. Also, on the third postnatal day, a high density of collagen IV fibrils was detected in the ILM along the vitreoretinal border as well as the pigmented layer (arrow). The vitreous body (VB) stroma also showed weak reaction. (E and F). Scale bar 100 μ m.




Fig 2: Immunohistochemical analysis of collagen IV during retinal development in mouse. Data used in quantification 10→ (-), 20→ (+), 30→ (++) ,40→ (+++) and 50→ (++++).

Discussion

The presented study shows that although the inner limiting membrane was formed by embryonic day 14, collagen type IV could not be found in any parts of the developing retina. Previous studies have shown that at this time, some of the epithelial tissues express collagen IV and that basement membrane begins to appear at its former area. The question was: why were we not able to detect this protein? It could be assumed that although collagen IV should appear at this time, it is possible that its concentration in contrast to other proteins in the basement membrane during the days to follow is undetectably low. The other possibility is the kind of antibody used in this study. In this investigation we preferred to use a monoclonal antibody as it binds with only one specific epitope of collagen IV (in contrast to a polyclonal antibody which binds with several epitopes of a protein); therefore, it is possible that based on this very specific reaction, the number of active sites were not satisfactorily detectable. It could be possible that if we used a polyclonal antibody, the rate of reaction could be higher. The first immunostaining was weakly detected on the late 15th embryonic day. This labeling was detected in different eye parts such as the inner and outer limiting membranes, and the optic cup on day 17. The development of collagen type IV at this stage represents the special role of cell to cell interaction in retinal layers. Differentiation of these layers into cell types reflects an important role in retinal development (21). Observing the labeling reactions indicated that cell interactions continued in the following days. The cell interaction intensity gradually increased in a way that they con-

cluded at their highest possible degree on the 3rd day after birth. No remarkable change was recorded afterwards. Results of a similar study showed that the retinal development period is from 15.5 days of gestation to the 4th postnatal day (22). Also, developmental stages of rodents such as rats indicate that the final development of their visual organ and the differentiation of these layer of immature retinal cells from specialized ones occur in the last embryonic week particularly in the early stages of postnatal development (23). Investigations show that the retinal structure is completely formed from invagination of the two optic cup cell strata (24). The internal layer of the retina is covered with an inner limiting membrane which is adjacent to the vitreous body, whereas the outer limiting membrane is adjacent to the internal surface of the outer layer of the optic cup retina (25). The basement membrane formation is dependent on the existence of specific proteins among which collagen IV is the most predominant (26). Therefore, due to the fact that elimination of collagen type IV could result in deformation of the basement membrane, one can find the vital role of this type of protein in basement membrane formation (27). Besides, the basement membrane covering the internal layer of retinal cells (ganglionic cells) makes the adjacency of this layer to the developing vitreous body possible. The vitreous body, exchanging ions and molecules as well as developmental signals, helps in the differentiation of the basement membrane (28). This is of prime importance because some of the mutants and transgenic that can not expressed collagen have visual defect (29). The labeling of collagen in ECM of retinal cells, and particularly its high concentration in ganglionic cells showed that the ganglionic layer development is dependent on the inner limiting membrane and needs the collagen type IV producing ECM (30). Moreover, the outer limiting membrane, connecting the inner layer of the optic cup with the outer layer of the pigmented epithelium, guarantees extended cell interactions which help the formation and differentiation of rod and cone cells as well as the retinal pigment epithelium (31).

Conclusion

Therefore, any disorder in the development of collagen IV in the optic cup affects the BM in this area and may result in anomalies such as defects in the natural development of pigmented epithelium or lack of accurate differentiation

of rod and cone cells. Hence, it seems that retinal development depends on various molecular changes and different sugars and proteins such as laminin and fibronectin, as well as different kinds of collagen especially collagen type IV. In other words, the development of retina requires various outer cell compounds including the inner limiting membrane and the extra cellular matrix. These compounds are mainly dependent on their composing elements especially collagen type IV to develop and play an important part in differentiation of cell layer.

Acknowledgments

This work was supported by a research grant (number 87040) from offices of the Vice-Chancellor for Research Affairs at Zabol and Mashhad Universities of Medical Sciences. The authors wish to thank both Vice Chancellors for Research Affairs for their financial support. We also wish to thank Mrs. Motejadded for her excellent technical assistance. There is no conflict of interest in this article.

References

- Berkholtz CB, Lai BE, Woodruff TK, Shea LD. Distribution of extracellular matrix proteins type I collagen, type IV collagen, fibronectin, and laminin in mouse folliculogenesis. *Histochem Cell Biol.* 2006; 126(5): 583-592.
- Halfter W, Dong S, Schurer B, Osanger A, Schneider W, Ruegg M, et al. Composition, synthesis, and assembly of the embryonic chick retinal basal lamina. *Dev Biol.* 2000; 220(2): 111-128.
- Halfter W, Dong S, Dong A, Eller AW, Nischt R. Origin and turnover of ECM proteins from the inner limiting membrane and vitreous body. *Eye.* 2008; 22(10): 1207-1213.
- Aisenbrey S, Zhang M, Bacher D, Yee J, Brunken WJ, Hunter DD. Retinal pigment epithelial cells synthesize laminins, including laminin 5, and adhere to them through alpha3- and alpha6-containing integrins. *Invest Ophthalmol Vis Sci.* 2006; 47(12): 5537-5544.
- Lin WL. Immunogold localization of extracellular matrix molecules in Bruch's membrane of the rat. *Curr Eye Res.* 1989; 8(11): 1171-1178.
- Halfter W, Dong S, Balasubramani M, Bier ME. Temporary disruption of the retinal basal lamina and its effect on retinal histogenesis. *Dev Biol.* 2001; 238(1): 79-96.
- Snead DR, Cullen N, James S, Poulson AV, Morris AH, Lukaris A, et al. Hyperconvolution of the inner limiting membrane in vitreomaculopathies. *Graefes Arch Clin Exp Ophthalmol.* 2004; 42(10): 853-862.
- Halfter W, Von Boxberg Y. Axonal growth on solubilized and reconstituted matrix from the embryonic chicken retina Inner limiting membrane. *Eur J Neurosci.* 1992; 4(9): 840-852.
- Miceli MV, Newsome DA. Effects of extracellular matrix and Bruch's membrane on retinal outer segment phagocytosis by cultured human retinal pigment epithelium. *Curr Eye Res.* 1996; 15(1): 17-26.
- Ishizaki M, Westerhausen-Larson A, Kino J, Hayashi T, Kao WW. Distribution of collagen IV in human ocular tissues. *Invest Ophthalmol Vis Sci.* 1993; 34(9): 2680-2689.
- Sarthy V. Collagen IV mRNA expression during development of the mouse retina: an in situ hybridization study. *Invest Ophthalmol Vis Sci.* 1993; 34(1): 145-152.
- Jo N, Ju M, Nishijima K, Robinson GS, Adamis AP, Shima DT, et al. Inhibitory effect of an antibody to cryptic collagen type IV epitopes on choroidal neovascularization. *Mol Vis.* 2006; 12: 1243-1249.
- Kern P, Laurent M, Lim A, Regnault F, Courtois Y. Interaction of bovine epithelial lens (BEL) cells with extracellular matrix (ECM) and eye-derived growth factor (EDGF). II. Partial re-expression of the differentiated collagen distribution and phenotype. *Exp Cell Res.* 1983; 149(1): 85-93.
- Ponsioen TL, van Luyn MJ, van der Worp RJ, van Meurs JC, Hooymans JM, Los LI. Collagen distribution in the human vitreoretinal interface. *Invest Ophthalmol Vis Sci.* 2008; 49(9): 4089-4095.
- Tsilibary EC. Microvascular basement membranes in diabetes mellitus. *J Pathol.* 2003; 200(4): 537-546.
- Bradshaw AD, McNagny KM, Gervin DB, Cann GM, Graf T, Clegg DO. Integrin alpha 2 beta 1 mediates interactions between developing embryonic retinal cells and collagen. *Development.* 1995; 121(11): 3593-3602.
- Boot-Handford RP, Kurkinen M, Prockop DJ. Steady-state levels of mRNAs coding for the type IV collagen and laminin olypeptide chains of basement membranes exhibit marked tissue-specific stoichiometric variations in the rat. *J Biol Chem.* 1987; 262(26): 12475-12478.
- Uemura A, Kusuhara S, Wiegand SJ, Yu RT, Nishikawa S. Tlx acts as a proangiogenic switch by regulating extracellular assembly of fibronectin matrices in retinal astrocytes. *J Clin Invest.* 2006; 116(2): 369-377.
- Das A, Frank RN, Zhang NL, Turczyn TJ. Ultrastructural localization of extracellular matrix components in human retinal vessels and Bruch's membrane. *Arch Ophthalmol.* 1990; 108(3): 421-429.
- Scheiffarth OF, Kampik A, Günther H, von der Mark K. Proteins of the extracellular matrix in vitreoretinal membranes. *Graefes Arch Clin Exp Ophthalmol.* 1988; 226(4): 357-361.
- Okada M, Ogino N, Matsumura M, Honda Y, Nagai Y. Histological and immunohistochemical study of idiopathic epiretinal membrane. *Ophthalmic Res.* 1995; 27(2): 118-128.
- Bai X, Dilworth DJ, Weng YC, Gould DB. Developmental distribution of collagen IV isoforms and relevance to ocular diseases. *Matrix Biol.* 2009; 28: 194-201.
- Connolly SE, Hores TA, Smith LE, D'Amore PA. Characterization of vascular development in the mouse retina. *Microvasc Res.* 1988; 36(3): 275-290.
- Canfield AE, Schor AM, Schor SL, Grant ME. The biosynthesis of extracellular-matrix components by bovine retinal endothelial cells displaying distinctive morphological phenotypes. *Biochem J.* 1986; 235(2): 375-383.
- Ito M, Nakashima M, Tsuchida N, Imaki J, Yoshioka M. Histogenesis of the intravitreal membrane and secondary vitreous in the mouse. *Invest Ophthalmol Vis Sci.* 2007; 48(5): 1923-1930.
- Kalebic T, Garbisa S, Glaser B, Liotta LA. Basement membrane collagen: degradation by migrating endothelial cells. *Science.* 1983; 221(4607): 281-83.
- Cagliero E, Maiello M, Boeri D, Roy S, Lorenzi M. Increased expression of basement membrane components in human endothelial cells cultured in high glucose. *J Clin Invest.* 1988; 82(2): 735-738.
- Snead DR, James S, Snead MP. Pathological changes in the vitreoretinal junction 1: epiretinal membrane formation. *Eye.* 2008; 22(10): 1310-1317.
- Frau E, Magnon C, Opolon P, Connault E, Opolon D, Beermann F, et al. A gene transfer comparative study of

Collagen IV Expression during Retinal Development

- HSA-conjugated antiangiogenic factors in a transgenic mouse model of metastatic ocular cancer. *Cancer Gene Ther.* 2007; 14(3): 251-261.
30. Snead MP, Snead DR, Richards AJ, Harrison JB, Poulson AV, Morris AH, et al. Clinical, histological and ultrastructural studies of the posterior hyaloid membrane. *Eye.* 2002; 16(4): 447-453.
31. Schmut O. The organization of tissues of the eye by different collagen types. *Albrecht Von Graefes Arch Klin Exp Ophthalmol.* 1978; 207(3): 189-199.
-