

# Effect of Mechanical Micro-Vibrations on The Efficiency of Leopard Inter-Species Somatic Cell Nuclear Transfer

Maryam Shahverdi, D.V.M.<sup>1,2</sup>, Vahid Akbarinejad, Ph.D.<sup>2\*</sup>, Azam Dalman, Ph.D.<sup>1</sup>, Mostafa Hajinasrollah, Ph.D.<sup>3</sup>, Mehdi Vojgani, Ph.D.<sup>2</sup>, Nima Tanhaei Vash, B.Sc.<sup>4</sup>, Mohammad Hossein Nasr-Esfahani, Ph.D.<sup>4</sup>, Poopak Eftekhari-Yazdi, Ph.D.<sup>1\*</sup>

1. Department of Embryology, Reproductive Biomedicine Research Centre, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

2. Department of Theriogenology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

3. Animal Core Facility, Reproductive Biomedicine Research Center, Royan Institute for Biotechnology, ACECR, Tehran, Iran

4. Department of Animal Biotechnology, Reproductive Biomedicine Research Center, Royan Institute for Biotechnology, ACECR, Isfahan, Iran

\*Corresponding Addresses: P.O.Box: 141556453, Department of Theriogenology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran  
P.O.Box: 16635-148, Department of Embryology, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran  
Emails: v\_akbarinejad@ut.ac.ir, eftekhari@royaninstitute.org

Received: Received: 29/January/2022, Accepted: 25/April/2022

## Abstract

**Objective:** Scarcity of oocytes for assisted reproduction in endangered species can be bypassed by interspecies somatic cell nuclear transfer (iSCNT). In Felids, domestic cat (*Felis catus*) oocytes can serve as recipients for the nucleus of the endangered Persian leopard (*Panthera pardus saxicolor*). However, *in vitro* oocyte maturation is still suboptimal in cats, whereas it has been reported to benefit from micro-vibration in non-felid species. Therefore, the present study is aimed to determine whether micro-vibration, applied during *in vitro* maturation (IVM), improves the embryogenic potential of cat oocytes transplanted with fibroblast nuclei of the Persian leopard.

**Materials and Methods:** In the experimental study, cat cumulus-oocyte complexes (COCs) were randomly assigned to the treatment group (micro-vibration) or control group (static culture). Resultant metaphase II (MII) oocytes were enucleated and reconstructed with nucleus transplants from leopard fibroblasts, followed by artificial oocyte activation and embryo culture under the same condition (static) for 7 days.

**Results:** While cumulus cell expansion and oocyte maturation profited from micro-vibration ( $P < 0.05$ ), the quantity and quality of blastocysts were significantly lower in micro-vibration than in the control group ( $P < 0.05$ ). The total number of blastocyst cells tended to be lower in the micro-vibration than in the control group ( $P = 0.075$ ). Nevertheless, the proportion of ICM and TE cells did not differ between the micro-vibration and control groups ( $P > 0.05$ ).

**Conclusion:** The present study indicated that micro-vibration at a frequency of 44 Hz for 5 secs per hour enhanced nuclear maturation and cumulus cell expansion of cat oocytes. However, exposure to micro-vibration during IVM impaired the survival rate of reconstructed oocytes during the iSCNT process and their developmental competence toward the blastocyst stage.

**Keywords:** Domestic Cat, *In Vitro* Maturation, *Panthera Pardus Saxicolor*, Somatic Cell Nuclear Transfer, Vibration

Cell Journal (Yakhteh), Vol 24, No 10, October 2022, Pages: 612-619

**Citation:** Shahverdi M, Akbarinejad V, Dalman A, Hajinasrollah M, Vojgani M, Tanhaei Vash N, Nasr-Esfahani MH, Eftekhari-Yazdi P. Effect of mechanical micro-vibrations on the efficiency of leopard inter-species somatic cell nuclear transfer. Cell J. 2022; 24(10): 612-619. doi: 10.22074/cellj.2022.8425.

This open-access article has been published under the terms of the Creative Commons Attribution Non-Commercial 3.0 (CC BY-NC 3.0).

## Introduction

According to the red list of the international union for conservation of nature, the population of the wild members of the Felidae family has dramatically dwindled over the past decades and all of them are presumed to be in jeopardy of extinction (1). This phenomenon has been attributed to various issues, including genetic, demographic, environmental, and human-associated factors restricting wild felids territory and rendering wildlife habitats uninhabitable (2, 3). In this regard, the Persian leopard (*Panthera pardus saxicolor*), which is one of the eight subspecies of leopard species (*Panthera pardus*), is mostly distributed in Iran and considered endangered; hence, protection of this subspecies is of utmost importance (4).

In the context of wildlife conservation, assisted

reproductive technologies can be applied to protect endangered animals (5) and there are numerous successful reports in leopard cats (6), African lions (7), and black-footed cats (8). In this regard, interspecies somatic cell nuclear transfer (iSCNT), in which the nucleus of one species is transferred to the enucleated oocyte of another species, could serve as a promising technique, particularly in wild cats affected by oocyte scarcity (9). Indeed, iSCNT using domestic cat oocytes has been applied to rescue a number of endangered feline species such as the African wild cat (10), sand cat (11), tiger (12), and cheetah (13). In these cases, the domestic cat oocyte was utilized as the recipient for the nucleus since it is the only feline species which is not considered endangered (10, 13). Nevertheless, iSCNT technique in feline species suffers from limitations. Chief among them is the fact

that despite major progress over recent years, rates of *in vitro* maturation (IVM) of domestic cat oocytes are still unsatisfactory for large-scale applications in iSCNT (14).

To improve IVM rate in cats, most studies have relied on chemical modifications of culture media, for instance, incorporation of growth factors or antioxidants (15). However, the possibilities are not exhausted with chemistry, and physical parameters are an option as well. Dynamic *in vitro* culture (IVC) using micro-vibration has been proposed to resemble *in vivo* conditions more closely, supporting improved oocyte maturation and development of resultant embryos (16-20). In validation of this notion, evaluating the effect of micro-vibration during IVM and IVC of embryos in pig, Mizobe et al. (17) observed that application of micro-vibration during IVM enhanced development of embryos up to the blastocyst stage, irrespective of the micro-vibration treatment during IVC, and the beneficial effect of micro-vibration was more pronounced during IVM than IVC. Also, micro-vibration has shown positive effects on embryo development in mice, cattle, pigs, and humans in the frequency range of 20-44 Hz.

In this study, we hypothesized that micro-vibration during IVM would result in cat oocytes that are more supportive of iSCNT. We conducted the present study to test this hypothesis, using oocytes from the domestic cat as recipients and fibroblasts from the Persian leopard as nucleus donors.

## Materials and Methods

All procedures were performed according to institutional guidelines on animal experimentation and care and approved by the Ethical Committee at Royan Institute (IR.ACECR.ROYAN.REC.1397.188). Unless otherwise mentioned, all materials used in this study were obtained from Sigma-Aldrich (St. Louis, MO, USA) and Gibco (Grand Island, NY, USA).

### Oocyte collection and IVM


Ovaries were collected from queens following routine ovariohysterectomy in veterinary clinics and kept in phosphate-buffered saline (PBS, Sigma-Aldrich, USA) at 4°C during transportation to the laboratory. After transportation, ovaries were washed (3 times) using PBS supplemented with 100 IU/ml penicillin and 100 mg/ml streptomycin at room temperature. Each ovary was sliced by a scalpel blade to release cumulus-oocyte complexes (COCs) into a washing medium (WM) containing HEPES tissue culture medium 199 (HTCM 199) supplemented with 10% fetal bovine serum (FBS, Gibco, Ireland). COCs with good morphology and quality (uniform in morphology with intact and dark ooplasm and more than three layers of cumulus cells) were selected for IVM (13). Maturation medium consisted of tissue culture medium

199 (TCM 199) supplemented with 2.5 mM sodium pyruvate, 1 ng/ml estradiol, 1 mM L-glutamine, 3% bovine serum albumin (BSA, Sigma-Aldrich, USA), 100 IU/ml penicillin and 100 mg/ml streptomycin, 25 ng/ml EGF, 100 ng/ml IGF1. The selected COCs were cultured in IVM medium droplets (5 to 10 COCs in each 50  $\mu$ L droplet), which were placed on tissue culture dishes and covered with mineral oil, at 38°C in 5% CO<sub>2</sub> for 24 hours. In the control (static) group, COCs received no more treatment, whereas, in the treatment (dynamic) group, COCs were subjected to mechanical micro-vibration at the frequency of 44 Hz for 5 seconds per hour using Vibroduct 1500 (SimSoTec GmbH, Cologne, Germany) (17, 18). Following IVM, cumulus expansion area was measured in 12 COCs using Fiji software (National Institutes of Health, Bethesda, MD, USA) as an image processing program, in which system units were changed to  $\mu$ m<sup>2</sup> by calibration. Afterward, cumulus cells were removed by treatment with hyaluronidase (300 IU/ml in HTCM199 containing 10% FBS) for 3 minutes, resulting in denuded oocytes. Oocyte maturation to metaphase II (MII) was scored by the presence of a polar body using a phase-contrast inverted microscope (Olympus CKX41, Tokyo, Japan), and the MII stage was confirmed by Hoechst staining (0.5 mg/ml) of 5 oocytes randomly selected from each group of denuded oocytes. Subsequently, the matured oocytes were deprived of the zona pellucida by incubation in pronase (2.5% in HTCM199 also containing 10% FBS) for 30 seconds on a warm plate and then allowed to recover for 15 minutes in 20% FBS.

### Isolation and preparation of leopard fibroblast cell line

The skin biopsy from the Persian leopard was provided to the Royan Institution by Tehran Zoo in April 2018. This step was implemented based on the methodology described previously by Dalman et al. (21). In brief, a biopsy from the ear of a Persian leopard was punched. After washing three times with PBS containing antibiotics (penicillin G 250 U/ml and streptomycin 250 mg/ml), the dermis was dissected into small pieces. After washing, the explants were cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 15% FBS, 1% penicillin and streptomycin, 2 mM L-glutamine, and 1 mM sodium pyruvate at 38°C in a 5% CO<sub>2</sub> incubator. After 10 days of culture when the cells reached confluency (80-90%), tissue explants were removed and cells were dissociated in trypsin (0.25% trypsin/EDTA) and centrifuged at 1200 rpm for 5 minutes and seeded. Early cell passages were cryopreserved in a medium containing 10% Dimethyl sulfoxide (DMSO, Sigma-Aldrich, USA) and stored in liquid nitrogen. For use in iSCNT experiments, cryopreserved fibroblasts were thawed, passaged 3 to 5 times in DMEM supplemented with 15% FBS at 38°C in a 5% CO<sub>2</sub> incubator, and

arrested in G0 by culturing them 48 hours in DMEM with 0.5% FBS (Fig.1).



**Fig.1:** Cultured leopard fibroblast cells (scale bar: 200  $\mu\text{m}$ ).

### Enucleation of oocytes

Enucleation was performed manually using a pulled Pasteur pipette as described previously by Hosseini et al. (22). In brief, zona-free oocytes were cultured in TCM199 supplemented with 0.41  $\mu\text{g/ml}$  demecolcine, 10% FBS, 10 mg/ml BSA, and 3 mg/ml polyvinyl alcohol (PVA) for 1.5 to 2 hours to induce cytoplasmic protrusion. Thereafter, manual enucleation of the oocyte was implemented under a stereomicroscope by a pulled Pasteur pipette. Before nuclear transfer, enucleated oocytes were maintained for 30 minutes in TCM199 supplemented with 10% FBS, 10 mg/ml BSA, and 3 mg/ml PVA.

### Interspecies somatic cell nuclear transfer

To place leopard cells in the vicinity of an enucleated oocyte, the population of fibroblast cells (about 500 cells) was transferred to 50  $\mu\text{l}$  drops of HTC199 supplemented with 10 mg/ml Phytohemagglutinin (PHA-P). Each enucleated oocyte was dropped on top of an intact and viable fibroblast and the two cells were pushed against each other with a Pasteur pipette. The resultant aggregates were placed in drops of HTC199 supplemented with 1 mg/ml PVA before fusion. The electrodes of the fusion chamber, lying 0.5 mm apart, were overlaid with a fusion medium consisting of TCM199, 0.3 mM mannitol, and 0.05% BSA. The aggregates were aligned manually and the cell membranes were fused by applying two pulses of direct electric current (1.75 kV/cm for 80 seconds and 1-second delay, Cryologic, Australia). Resultant reconstructed oocytes were maintained in HTC199 supplemented with 3 mg/ml PVA for 30 to 40 minutes to assess fusion success, before proceeding to activation.

### Artificial oocyte activation and *in vitro* embryo culture

Reconstructed oocytes were incubated for 5 minutes in

drops of activation medium composed of HTC199, 1 mg/ml BSA, and 10  $\mu\text{g/ml}$  calcium ionophore A23187. After brief washing in a drop of TCM199 supplemented with 10 mg/ml BSA and 1 mg/ml PVA, reconstructed oocytes were incubated 5 hours in 6-Dimethylamino purine (6-DMAP, 5  $\mu\text{g/ml}$ ). Given that zona-free oocytes were used for iSCNT; we chose a ‘well of well’ (WOW) culture system so as to prevent the embryos from sticking to each other while allowing for diffusion of embryo-derived growth factors between embryos (23). Six reconstructed oocytes were co-cultured in each WOW overlaid with 20  $\mu\text{l}$  of synthetic oviductal fluid (SOF) medium with oil on top, at 39°C, under an atmosphere of 5%  $\text{O}_2$ , 5%  $\text{CO}_2$ , and maximum humidity. *In vitro* culture conditions were identical between control and micro-vibration groups.

### Differential staining of embryos

Differential staining was used to identify inner cell mass (ICM) and trophectoderm (TE) cells by means of two DNA-specific fluorochromes, namely, propidium iodide (PI) and Hoechst 33342 (24). To this end, day-7 blastocysts were transferred to 2.5% Triton X100 in HTC199 plus 5 mg/ml BSA for 15 seconds, and then, they were transferred into HTC199-BSA medium containing 30  $\mu\text{g/ml}$  PI for 45 seconds and washed in HTC199-BSA medium. Afterward, blastocysts were transferred into 800  $\mu\text{l}$  of Hoechst 33342 (10  $\mu\text{g/ml}$  in Ethanol) for 15 minutes and were further mounted on glass slides and examined under a fluorescence microscope (Nikon Eclipse 50i, Tokyo, Japan). Counting of cell nuclei was performed on microphotographs taken with a digital camera. Following enumeration of cells in ICM and TE, the proportion of ICM and TE cells relative to total blastocyst cells (ICM+TE cells) were calculated for further analysis.

### Statistical analysis



There were three biological replicates in each experimental group. Continuous data, including COC expansion (size) and the number of blastocyst cells, were initially tested for normality using the Kolmogorov-Smirnov test, and as they had normal distribution, they were analyzed by t test. Non-continuous data, including rates (of IVM, reconstructed oocytes formation, degeneration, cleavage, and blastocyst formation) as well as proportions (of ICM and TE cells) were analyzed using logistic regression (GENMOD procedure), which generated odds ratios (ORs), as the strength of the difference between groups, and 95% confidence intervals (95% CIs). All analyses were conducted in SAS version 9.4 (SAS, 2013). Data are presented as mean  $\pm$  SEM. Differences at  $P < 0.05$  were considered statistically significant.

## Results



### The effect of micro-vibration on domestic cat oocyte IVM

Based on the projection area of the COCs (Fig.2A), micro-vibration during IVM supported greater expansion of the cumulus cells ( $9.48\text{E}+04 \pm 1.37\text{E}+04 \mu\text{m}^2$ ,  $n=12$ )

compared to the control group ( $4.74E+04 \pm 0.18E+04 \mu\text{m}^2$ ,  $n=12$ ) ( $P=0.0007$ , Fig.2B). Based on the presence of a polar body (Fig.3A), a higher proportion of cat oocytes progressed to MII in the micro-vibration group ( $91.70 \pm 1.77\%$ ,  $n=383$ ) compared to the control group ( $84.95 \pm 2.85\%$ ,  $n=383$ ,  $OR=1.816$ ,  $95\% CI=1.152-2.865$ ,  $P=0.010$ , Fig.3B).



**Fig.2:** Expansion of cumulus cells. **A.** Expansion of cumulus cells was highlighted by the dashed circles in control and micro-vibration groups (scale bar:  $100 \mu\text{m}$ ). **B.** Comparison of cumulus cell expansion in control ( $n=12$ ) and micro-vibration ( $n=12$ ) groups. \*;  $P<0.05$ .



**Fig.3:** Oocytes maturation. **A.** Extrusion of the first polar body was substantiated by Hoechst staining in a subset of MII-stage oocytes (five oocytes per replicate) (scale bar:  $20 \mu\text{m}$ ). **B.** Maturation rate of domestic cat oocytes subjected to *in vitro* maturation (IVM) under control ( $n=383$ ) and micro-vibration ( $n=383$ ) conditions. \*;  $P<0.05$ .

### The effect of micro-vibration on iSCNT rate and embryo development


Rate of reconstructed oocyte formation was less in the micro-vibration group ( $n=350$ ,  $23.92 \pm 5.82\%$ ) compared to the control group ( $n=327$ ,  $48.53 \pm 4.90\%$ ,  $OR=0.334$ ,  $95\% CI=0.240-0.465$ ,  $P<0.0001$ , Fig.4). Following activation, similar proportions of reconstructed oocytes degenerated in the two groups ( $11.02 \pm 0.26\%$ ,  $n=81$ , and  $15.81 \pm 5.72\%$ ,  $n=15$ , in the micro-vibration and control groups, respectively). Likewise, cleavage rates were similar in the two groups ( $83.19 \pm 0.44\%$ ,  $n=72$ , and  $90.63 \pm 1.73\%$ ,  $n=130$ ; in the micro-vibration and control groups, respectively) ( $P>0.10$ , Fig.4). However, the proportion of cleaved embryos that developed to the blastocyst stage after seven days of culture was lower in the micro-vibration ( $n=60$ ,  $6.61 \pm 0.82\%$ ) than in the control group ( $18.70 \pm 0.72\%$ ,  $n=118$ ,  $OR=0.312$ ,  $95\%$

CI=0.102-0.951, P=0.041, Fig.4).


**The effect of micro-vibration on iSCNT blastocysts quality**

Based on the absolute cell counts after differential staining (3 blastocysts in each group) (Fig.5A), the number of ICM cells was lower in the micro-vibration (19.67 ± 1.20) than in the control (25.33 ± 0.88) group

(P=0.019, Fig.5B), while the number of TE cells (70.00 ± 5.77, 86.33 ± 5.24) and total blastocyst cells (89.67 ± 6.94, 111.87 ± 6.01) were not significantly different in the micro-vibration and control groups, respectively (P>0.05, Fig.5B). According to the differential staining method, the ICM cell number did not differ between micro-vibration and control groups (22.00 ± 0.54% and 22.75 ± 0.66%, respectively), so did the TE cell number (78.00 ± 0.54% vs. 77.25 ± 0.66 %) (P>0.05, Fig.5B).



**Fig.4:** Rate of reconstructed oocyte formation, degeneration rate of reconstructed oocytes by day three of culture, cleavage rate of intact reconstructed oocytes, and rate of blastocyst formation relative to cleaved embryos in control and micro-vibration groups. \*; P<0.05.



**Fig.5:** Interspecies somatic cell nuclear transfer (iSCNT) blastocysts. **A.** Differential staining of iSCNT-generated blastocysts in control and micro-vibration groups (scale bar: 20 μm). **B.** Number of embryonic cells and proportion of inner cell mass (ICM) and trophectoderm (TE) cells relative to the total number of blastocyst cells in the control (n=3) and micro-vibration (n=3) groups. \*; P<0.05.

## Discussion

An inadequate rate of IVM in the domestic cat impedes the success of iSCNT in wild felids (17, 18). The present study aimed to investigate the effect of micro-vibration, applied during IVM, on the maturation of domestic cat oocytes and their embryogenic potential after iSCNT from Persian leopard fibroblasts. The results showed that to apply micro-vibration with the frequency of 44 Hz for 5 seconds per hour, the maturation rate of oocytes was improved but the blastocysts formation rate was decreased.

Our results show that micro-vibration enhanced the expansion of COCs as well as the MII progression of oocytes in the domestic cat. This is consistent with reports in pigs (25, 28). In particular, Mizobe et al. (17) demonstrated a greater extent of cumulus cell expansion in porcine COCs exposed to mechanical micro-vibration during IVM, although these authors did not observe any significant influence of micro-vibration on the proportion of oocytes reaching the MII stage. Mechanical micro-vibration has also proven itself in somatic cell applications, augmenting the proliferation, differentiation, and secretion parameters (27-29). Since COCs are comprised of somatic cells encasing an oocyte, it seems logical that micro-vibration could enhance oocyte maturation through facilitation of expansion in cumulus cells, which govern the maturation and developmental competence of oocytes (30). As an expected result of the enhancement, the IVM oocytes could be better prepared for their biological tasks, including fertilization and embryonic development (31).

Contrary to our expectations, cat oocytes subjected to micro-vibration during IVM were more vulnerable to iSCNT procedure, as measured by rates of reconstructed oocyte formation, which were inferior in the micro-vibration group compared to the control group. This observation implies that despite improved cumulus cell expansion and oocyte maturation to MII, the oocytes were probably damaged in other ways, thereby detracting from their ability to survive the subsequent procedures. Since the settings of intensity, duration, and frequency of micro-vibration used in this study were adopted from studies in other species (e.g. pig (17), human (18), mice (19), and cows (20)), it is conceivable that the settings did not suit feline oocytes, reducing their ability to support iSCNT in terms of embryo development (17,18). In this context, we speculate that the plasma membrane may be key to understanding what happened in our study. The plasma membrane is the outermost cellular component that receives the shear stress induced by micro-vibration, and plasma membrane also plays a critical role as a signal transduction interface between external and intracellular environment, subject to modulation by its structural and biochemical properties, particularly its lipid composition and organization (32-34). It is of note that lipid composition and organization determine plasma membrane integrity, prerequisite for cellular homeostasis, and tolerance to extracellular pressures and manipulations (32, 33, 35).

Hence, the adverse effects of micro-vibration settings - adopted from other species - on cat oocyte survival during iSCNT could be attributed to the difference(s) between cat and other species in terms of oolemma's physical and biochemical properties, including lipid composition (36).

After activation of the reconstructed oocytes, degeneration and cleavage rate did not differ between the two experimental groups, while the micro-vibration group produced fewer embryos with diminished proliferation. This negative effect of micro-vibration on embryonic development may have resulted from settings (e.g. duration and/or intensity of micro-vibration) which were inappropriate for the domestic cat oocyte. In this regard, while evaluating the effect of various durations of micro-vibration on IVM of porcine oocytes, Mizobe et al. (17) observed that prolonged exposure to mechanical micro-vibration reduced parthenogenetic development of oocytes up to the blastocyst stage. Moreover, aiming to ascertain the optimum frequency of micro-vibration for IVM and IVC of bovine oocytes and embryos, Takahashi et al. (20) found that an excessive micro-vibration intensity reduced development to the blastocyst stage as well as blastocyst cell proliferation. In the present study, the nuclear maturation rate was improved by micro-vibration, but cytoplasmic maturation remains to be assessed under static vs dynamic conditions. Therefore, the adverse effect of micro-vibration on oocyte developmental competence in the current study could have been mediated through the disruption of processes involved in oocyte cytoplasmic maturation. Indeed, normal development relies on coordinated nuclear and cytoplasmic maturation of oocytes (37, 38). Interference of micro-vibration with the oocytes' cytoplasmic maturation during IVM might have led to disruption of molecular mechanisms regulating proliferation of the future embryonic cells. Among these mechanisms, we envision, for example, the tissue growth factor  $\beta$  superfamily proteins (39) and the Hippo signaling pathway (40).

In this study, the most important limitation that we faced was the scarcity of cat ovaries and oocytes and it restricted the study from finding the optimal frequency for cat oocytes and iSCNT embryos.

## Conclusion

Micro-vibration at the frequency of 44 Hz for 5 seconds per hour during IVM augmented cumulus cell expansion and meiotic maturation in domestic cat oocytes. However, the oocytes that matured under the influence of micro-vibration were less tolerant of iSCNT and produced lower numbers of reconstructed oocytes. In addition, these reconstructed oocytes were less capable of developing into blastocysts, which also contained fewer cells. Therefore, it appears that this field of research requires further studies to tailor the characteristics of micro-vibration during IVM of domestic cat oocytes for use in either intraspecies or interspecies purposes. In particular, it should be clarified if the low tolerance of micro-vibrated cat oocytes to iSCNT stemmed from the micro-vibration, or from the

specific settings used (intensity, duration, and frequency of micro-vibration). It would also be helpful to examine the expression of pluripotency and trophectoderm markers in the blastocysts like *OCT4*, *CDX2*.

## Acknowledgments

We express gratitude to the staff of the cloning laboratory at Royan Biotechnology Institute and the Embryology Laboratory at Royan Institute. This study was supported by Royan institute (grant number 97000012); and the Faculty of Veterinary Medicine, University of Tehran (grant number 30854/6/3). The authors declare that there is no conflict of interest regarding the publication of this paper.

## Authors' Contributions

M.Sh., A.D.; Participated in the investigation, methodology, writing, review, and editing. V.A., P.E.-Y.; Funding acquisition, project administration, supervision, formal analysis, review, and editing. M.H.; Prepared cat's ovaries following ovariectomy. M.V., M.H.N-E.; Provided critical revision of the study. N.T.V.; Contributed to all experimental work and data. All authors read and approved the final manuscript.

## References

- Wildt D, Phillips L, Simmons L, Chakraborty P, Brown J, Howard J, et al. A comparative analysis of ejaculate and hormonal characteristics of the captive male cheetah, tiger, leopard and puma. *Biol Reprod*. 1988; 38: 245-255.
- Jacobson AP, Gerngross P, Lemeris JR Jr, Schoonover RF, Anco C, Breitenmoser-Würsten C, et al. Leopard (*Panthera pardus*) status, distribution, and the research efforts across its range. *PeerJ*. 2016; 4: 1-28.
- Thuwanut P, Tipkantha W, Siriaroonrat B, Comizzoli B, Chatdarong K. Beneficial effect of extracellular adenosine 5' triphosphate treatment on the Indochinese leopard (*Panthera pardus delacouri*) sperm quality after cryopreservation. *Reprod Domest Anim*. 2017; 52(52): 269-274.
- Naderi M, Farashi A, Erdi MA. Persian leopard's (*Panthera pardus saxicolor*) unnatural mortality factors analysis in Iran. *PLoS One*. 2018; 13(4): 1-12.
- Mikołajewska N, Müller K, Nizański W, Jewgenow K. Vitrification of domestic cat oocytes—effect on viability and integrity of subcellular structures. *Reprod Domest Anim*. 2012; 47 Suppl 6: 295-299.
- Goodrowe KL, Miller AM, Wildt DE. In vitro fertilization of gonadotropin-stimulated leopard cat (*Felis bengdemis*) follicular oocytes. *J exp Zool*. 1989; 252(1): 89-95.
- Goeritz F, Painer J, Jewgenow K, Hermes R, Rasmussen K, Dehnhard M, et al. Embryo retrieval after hormonal treatment to control ovarian function and non-surgical artificial insemination in African lions (*Panthera leo*). *Reprod Domest Anim*. 2012; 47 Suppl 6: 156-160.
- Pope CE, Gómez MC, Galiguis J, Dresser BI. Applying embryo cryopreservation technologies to the production of domestic and black-footed cats. *Reprod Domest Anim*. 2012; 47 Suppl 6: 125-129.
- Cordova A, King WA, Mastromonaco GF. Choosing a culture medium for SCNT and iSCNT reconstructed embryos: from domestic to wildlife species. *J Anim Sci Technol*. 2017; 59: 24
- Gómez MC, Pope CE, Giraldo A, Lyons LA, Harris RF, King AL, et al. Birth of african wildcat cloned kittens born from domestic cats. *Cloning Stem Cells*. 2004; 6(3): 247-258.
- Gómez MC, Pope CE, Kutner RH, Ricks DM, Lyons LA, Ruhe M, et al. Nuclear transfer of sand cat cells into enucleated domestic cat oocytes is affected by cryopreservation of donor cells. *Cloning Stem Cells*. 2008; 10(4): 469-483.
- Moro LN, Jarazo J, Buemo C, Hiriart MI, Sestelo A, Salamone DF. Tiger, Bengal and Domestic cat embryos produced by homospesific and interspecific zona-free nuclear transfer. *Reprod Domest Anim*. 2015; 50(5): 849-857.
- Moro LN, Hiriart MI, Buemo C, Jarazo J, Sestelo A, Veraguas D, et al. Cheetah interspecific SCNT followed by embryo aggregation improves in vitro development but not pluripotent gene expression. *Reproduction*. 2015; 150(1): 1-10.
- Amamath D, Li X, Kato Y, Tsunoda Y. Gene expression in individual bovine somatic cell cloned embryos at the 8-cell and blastocyst stages of preimplantation development. *J Reprod Dev*. 2007; 53(6): 1247-1263.
- Ochota M, Pasięka A, Nizański W. Superoxide dismutase and taurine supplementation improves in vitro blastocyst yield from poor-quality feline oocytes. *Theriogenology*. 2016; 85(5): 922-927.
- Fauci LJ, Dillon R. Biofluidmechanics of reproduction. *Annu Rev Fluid Mech*. 2006; 38: 371-394.
- Mizobe Y, Yoshida M, Miyoshi K. Enhancement of cytoplasmic maturation of in vitro matured pig oocytes by mechanical vibration. *J Reprod Dev*. 2010; 56(2): 285-290.
- Isachenko V, Maettner R, Sterzik K, Strehler E, Kreinberg R, Hancke K, et al. In-vitro culture of human embryos with mechanical micro-vibration increases implantation rates. *Reprod Bio Med Online*. 2011; 22(6): 536-544.
- Hur YS, Park JH, Ryu EK, Park SJ, Lee JH, Lee SH, et al. Effect of micro-vibration culture system on embryo development. *J Assist Reprod Genet*. 2013; 30(6): 835-841.
- Takahashi M, Honda T, Hatoya S, Inaba T, Kawate N, Tamada H. Efficacy of mechanical micro-vibration in the development of bovine embryos during in vitro maturation and culture. *J Vet Med Sci*. 2018; 80(3): 532-535.
- Dalman A, Eftekhari-Yazdi P, Valojerdi MR, Shahverdi A, Gourabi H, Janzamin E, et al. Synchronizing cell cycle of goat fibroblasts by serum starvation causes apoptosis. *Reprod Domest Anim*. 2010; 45(5): e46-e53.
- Hosseini SM, Hajian M, Moulavi F, Asgari V, Forouzanfar M, Nasr-Esfahani MH. Cloned sheep blastocysts derived from oocytes enucleated manually using a pulled pasteur pipette. *Cell Reprogram*. 2013; 15(1): 15-23.
- Vajta G, Peura TT, Holm P, Paldi A, Greve T, Trounson AO, et al. New method for culture of zona-included or zona-free embryos: the Well of the Well (WOW) system. *Mol Reprod Dev*. 2000; 55(3): 256-264.
- Papaioannou VE, Ebert KM. The preimplantation pig embryo: cell number allocation to trophectoderm and inner cell mass of the blastocyst in vivo and in vitro. *Development*. 1988; 102(4): 793-803.
- Comizzoli P, Wildt DE, Pukazhenthi BS. Overcoming poor in vitro nuclear maturation and developmental competence of domestic cat oocytes during the non-breeding season. *Reproduction*. 2003; 126(6): 809-816.
- Fernandez-Gonzalez L, Hribal R, Stagegaard J, Zahmel J, Jewgenow K. Production of lion (*Panthera leo*) blastocysts after in vitro maturation of oocytes and intracytoplasmic sperm injection. *Theriogenology*. 2015; 83(6): 995-999.
- Puig F, Rico F, Almendros I, Navajas D, Montserrat JM, Farre R. Vibration enhances interleukin-8 release in a cell model of snoring-induced airway inflammation. *Sleep*. 2005; 28(10): 1312-1316.
- Kaupp JA, Waldman SD. Mechanical vibrations increase the proliferation of articular chondrocytes in high-density culture. *Proc Inst Mech Eng H*. 2008; 222(5): 695-703.
- Ito Y, Kimura T, Nam K, Katoh A, Masuzawa T, Kishida A. Effects of vibration on differentiation of cultured PC12 cells. *Biotechnol Bioeng*. 2011; 108(3): 592-599.
- Dhali A, Javvaji PK, Kolte AP, Francis JR, Roy SC, Sejian V. Temporal expression of cumulus cell marker genes during in vitro maturation and oocyte developmental competence. *J Assist Reprod Genet*. 2017; 34(11): 1493-1500.
- De Matos DG, Miller K, Scott R, Tran CA, Kagan D, Nataraja SG, et al. Leukemia inhibitory factor induces cumulus expansion in immature human and mouse oocytes and improves mouse two-cell rate and delivery rates when it is present during mouse in vitro oocyte maturation. *Fertil Steril*. 2008; 90(6): 2367-2375.
- White CR, Frangos JA. The shear stress of it all: the cell membrane

- and mechanochemical transduction. *Philos Trans R Soc Lond B Biol Sci.* 2007; 362(1484): 1459-1467.
33. Le Roux AL, Quiroga X, Walani N, Arroyo M, Roca-Cusachs P. The plasma membrane as a mechanochemical transducer. *Philos Trans R Soc Lond B Biol Sci.* 2019; 374(1779): 1-15.
  34. Tozzi C, Walani N, Arroyo M. Out-of-equilibrium mechanochemistry and self-organization of fluid membranes interacting with curved proteins. *New J Phys.* 2019; 21: 1-23.
  35. McNeil PL, Steinhardt RA. Loss, restoration, and maintenance of plasma membrane integrity. *J Cell Biol.* 1997; 137(1): 1-4.
  36. Apparicio M, Ferreira CR, Tata A, Santos VG, Alves AE, Mostachio GQ, et al. Chemical composition of lipids present in cat and dog oocyte by matrix-assisted desorption ionization mass spectrometry (MALDI- MS). *Reprod Domest Anim.* 2012; 47 Suppl 6: 113-117.
  37. Watson AJ. Oocyte cytoplasmic maturation: a key mediator of oocyte and embryo developmental competence. *J Anim Sci.* 2007; 85(13 Suppl): E1-3.
  38. Conti M, Franciosi F. Acquisition of oocyte competence to develop as an embryo: integrated nuclear and cytoplasmic events. *Hum Reprod Update.* 2018; 24(3): 245-266.
  39. Anani S, Bhat S, Honma-Yamanaka N, Krawchuk D, Yamanaka, Y. Initiation of Hippo signaling is linked to polarity rather than to cell position in the pre-implantation mouse embryo. *Development.* 2014; 141(14): 2813-2824.
  40. Liu C, Peng G, Jing N. TGF- $\beta$  signaling pathway in early mouse development and embryonic stem cells. *Acta Biochim Biophys Sin (Shanghai).* 2018; 50(1): 68-73.
-