

Supplementary Information for

Cordycepin Suppresses The Malignant Phenotypes of Colon Cancer Cells through The GSK3 β / β -catenin/cyclin D1 Signaling Pathway

Jie Xu, B.Sc.¹, Xia Shen, B.Sc.², Daozhong Sun, B.Sc.¹, Yanjie Zhu, B.Sc.^{3*}

1. Department of General Surgery, Zhejiang Greentown Cardiovascular Hospital, Hangzhou, Zhejiang, China
2. Department of Emergency, Zhejiang Greentown Cardiovascular Hospital, Hangzhou, Zhejiang, China
3. Department of Dermatology, The Second People's Hospital of Yuhang District, Hangzhou, Zhejiang, China

*Corresponding Address: Department of Dermatology, The Second People's Hospital of Yuhang District, Hangzhou, Zhejiang, China
Email: 1048950853@qq.com




Fig.S1: Effects of cordycepin on the pulmonary metastasis of colon cancer cells. **A.** Cell viability of FHC cells was detected by MTT assay after treatment with different concentrations of cordycepin (control, 20 μ M, 40 μ M and 80 μ M). **B.** Lung metastasis of HT-29 cells *in vivo* was evaluated after the nude mice were injected with HT-29 cells treated with or without 80 μ M cordycepin for 24 hours. The ordinate indicates the number of nodules per section. *; P<0.05 and ***, P<0.001.