

Supplementary Information for

Study of The Correlation between miR-106a, miR-125b, and miR-330 on Multiple Sclerosis Patients by Targeting TNFSF4 and SP1 in NF- κ B/TNF- α Pathway: A Case-Control Study

Nasrin Hadi, M.Sc.¹, Seyed Morteza Seifati, Ph.D.¹, Behnaz Nateghi, M.Sc.^{2,3}, Parisa Ravaghi, M.Sc.⁴, Farinaz Khosravian, M.Sc.^{2,3}, Faezeh Namazi, M.Sc.¹, Maryam Fotouhi Firouzabad, M.Sc.¹, Vahid Shaygannejad, M.D.⁵, Mansoor Salehi, Ph.D.^{2,3,6*}

1. Medical Biotechnology Research Center, Ashkezar Branch, Islamic Azad University, Ashkezar, Yazd, Iran
2. Medical Genetics Research Center of Genome, Isfahan University of Medical Sciences, Isfahan, Iran
3. Cellular, Molecular, and Genetics Research Center, Isfahan University of Medical Sciences, Isfahan, Iran
4. Department of Biology, Faculty of Science, Shahid Chamran University of Ahvaz, Ahvaz, Iran
5. Department of Neurology, Isfahan Neurosciences Research Center, Isfahan University of Medical Sciences, Isfahan, Iran
6. Department of Genetics and Molecular Biology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

*Corresponding Address: P.O.Box: 81746-73461, Department of Genetics and Molecular Biology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran
Email: m_salehi@med.mui.ac.ir




Fig.S1: Gene enrichment analysis. Bar graphs show the significance of the P value in each pathway in which the related genes are involved in multiple sclerosis.