

Supplementary Information for

Establishment of A Three-Dimensional Culture Condition for The U266 Cell Line Based on Peripheral Blood Plasma-Derived Fibrin Gels

Mahshid Jomehpour, M.Sc.^{1#}, Mohammadmahdi Khosravi, M.Sc.^{1#}, Mehrnaz Janfada, M.Sc.^{1#}, Saeid Abroun, Ph.D.¹, Sadaf Vahdat, Ph.D.^{2*}

1. Department of Hematology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

2. Applied Cell Sciences Division, Department of Hematology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran




Fig.S1: The effect of different calcium chloride concentrations on the gelation of plasma. Numbers displayed on vials indicate different concentrations (0-5 mg/ml) of calcium chloride.




Fig.S2: The effect of administration of different concentrations of collagenase type I on the viability of U266 cells after fibrin gel digestion. The data is presented in the form of mean $\pm$ standard deviation.

Received: 02/October/2022, Revised 30/January/2023, Accepted: 14/February/2023

These authors equally contributed to this work.

*Corresponding Address: P.O.Box: 14115-111, Applied Cell Sciences Division, Department of Hematology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Email: s_vahdat@modares.ac.ir